

Staggered Chiral Perturbation Theory for Heavy-Light Mesons

C. Aubin

Dept. of Physics, Columbia University, New York, NY 10027

C. Bernard

Washington University, St. Louis, MO 63130

Abstract

We incorporate heavy-light mesons into staggered chiral perturbation theory (SXPT), working to leading order in $1/m_Q$, where m_Q is the heavy quark mass. At first non-trivial order in the chiral expansion, staggered taste violations affect the chiral logarithms for heavy-light quantities only through the light meson propagators in loops. There are also new analytic contributions coming from additional terms in the Lagrangian involving heavy-light and light mesons. Using this heavy-light SXPT, we perform the one-loop calculation of the B (or D) meson leptonic decay constant in the partially quenched and full QCD cases. In our treatment, we assume the validity both of the “fourth root trick” to reduce four staggered tastes to one, and of the SXPT prescription to represent this trick by insertions of factors of $1/4$ for each sea quark loop.

PACS numbers: 12.39.Fe, 12.39.Hg, 11.30.Rd, 12.38.Gc

I. INTRODUCTION

The lattice can make a major contribution to the understanding of flavor physics through the computation of the properties of heavy-light mesons. Among the important quantities to calculate are the heavy-light decay constants, form factors, and B parameters. (See Ref. [1, 2] for recent reviews.) However, the “traditional” lattice approach, using quite massive light quarks, is likely to produce results that have large systematic errors coming from the long chiral extrapolation [3]. An alternative approach that seems very promising is to use staggered (Kogut-Susskind) fermions for both the light valence and sea quarks [4, 5, 6, 7, 8, 9]. An exact chiral symmetry for staggered quarks at finite lattice spacing allows simulations to be performed at small quark mass, deep in the chiral regime.

Staggered quarks have “taste” symmetry, a four-fold remnant of the doubling symmetry. Taste symmetry is broken at $\mathcal{O}(a^2)$ in the lattice spacing, resulting in rather large discretization effects. Even at the smaller lattice spacing employed in current MILC simulations with improved staggered quarks [10] (the “fine” set with lattice spacing $a \approx 0.09$ fm), the effects are not negligible in any quantity which is sensitive to pseudoscalar meson loops. This has been understood in the light meson sector using staggered chiral perturbation theory (SXPT) [11, 12, 13]. We now extend this program to include heavy quarks, thus merging heavy quark effective theory (HQET) with SXPT. We then calculate the leptonic decay constants, f_B and f_{B_s} , at one loop in SXPT in both the partially quenched and full QCD cases. This is one of the simpler quantities to calculate both in lattice simulations [1, 2] and in the SXPT formalism.

Simulations with staggered fermions such as Refs. [4, 5, 6, 7, 8, 9, 10], use the “fourth root trick” [14], designed to reduce the number of tastes per flavor from 4 to 1 in the continuum limit. The validity of the fourth root trick has not been proven, although various recent studies have, in our opinion, made it rather plausible [15]. In the following, we *assume* that this trick is valid, and that it can be represented in the chiral theory by insertions of factors of $1/4$ for each sea quark loop [12, 13]. We employ a quark-flow analysis [16] to locate these sea quark loops. The “replica method” [17] is an alternative technique for inserting the appropriate factors. It is theoretically cleaner than quark flow, but usually somewhat more complicated to implement in practice. One would start with n_S staggered fields for each continuum sea-quark flavor desired, and take the limit $n_S \rightarrow 1/4$ at the end. The two

approaches yield identical results in the current case, and in all other cases that have been investigated to date.

A brief discussion of the current results, as well as the application to semileptonic form factors, was previously presented in Ref. [18]. A detailed description of the semileptonic calculation is in preparation [19]. The SXPT forms obtained have already been used for the chiral (and continuum) extrapolations of numerical lattice results for heavy-light semileptonic [8] and leptonic [9] decays.

In this paper the heavy quark mass m_Q is taken to be large compared to Λ_{QCD} , so that working to leading order in the heavy quark expansion (*i.e.*, neglecting $1/m_Q$ terms) is a reasonable first approximation. We also do not include, in our effective theory, discretization errors coming from the heavy quark. We assume that such errors can be adequately estimated independently, using HQET as the effective theory description of the lattice heavy quark [2, 20, 21]. However, in order to separate light and heavy quark discretization errors, our analysis requires that the heavy quark mass not be much greater than one in lattice units. The rather counterintuitive restriction arises because the heavy quark action no longer sufficiently suppresses doublers when $am_Q \gg 1$. Indeed, in the large am_Q limit, the heavy quark action becomes a static lattice action, where there is no suppression of spatial doublers at all, since a static quark’s energy is independent of its momentum. When heavy quark doublers are insufficiently suppressed, the light and heavy quarks in a heavy-light meson can exchange gluons of momenta π/a , leading to “mixed” 4-quark operators (products of heavy and light bilinears) in the Symanzik action that violate taste. In order to avoid treating this complication — which would inextricably mix light- and heavy-quark discretization errors — we do not allow $am_Q \gg 1$ in our analysis. On the other hand, typical values of am_Q in practical calculations with the Fermilab [22] or NRQCD [23] heavy quarks are likely to be acceptable. With this restriction, all taste-violations in the $\mathcal{O}(a^2)$ Symanzik action occur solely in the light quark sector; the taste-violating terms are thus the same as in Refs. [11, 13]. More details, including a discussion based on the symmetries of the lattice action, appear in Sec. II.

Incorporating heavy quarks into SXPT produces a large number of terms in the chiral Lagrangian involving combinations of the heavy-light and the light mesons. All of these terms, which arise due to the taste symmetry breaking, are however non-leading: They do not appear in one-loop diagrams for the heavy-light quantities and enter the decay constant

result through a single analytic term. Non-trivial taste violations arise only in the light meson propagators in the one-loop diagrams. Since the taste-violating low energy constants in the light meson sector have been determined from simulations and SXPT fits [10, 24, 25], the new analytic term is the only additional fit parameter beyond those low-energy constants that appear in the continuum, such as the B - B^* - π coupling g_π .

This paper is organized as follows. We first describe the Symanzik action to $\mathcal{O}(a^2)$ in Sec. II. The SXPT Lagrangian, here including heavy-light mesons, is then constructed from a spurion analysis in Secs. III and IV, where the latter focuses on the $\mathcal{O}(a^2)$ terms involving heavy-light mesons. In Sec. V we calculate the one-loop expression for the chiral logarithms that arise in the leptonic decay constant, f_{B_x} , where x is the flavor of the light valence quark. We write down the final results for various cases in Sec. VI and discuss some features of the low energy constants (LECs) in Sec. VII. Finite volume effects are treated in Sec. VIII. We finish with some remarks and conclusions in Sec. IX.

II. SYMANZIK ACTION

For concreteness, we consider here a heavy quark that is simulated either with NRQCD [23] or with the Fermilab interpretation [22] of a clover or more highly improved quark.¹ What is crucial for us about these possible heavy quark actions is that they describe a single physical fermion — doubler² masses are assumed to be larger than the physical heavy quark mass by an amount of order of the cutoff, *i.e.*, $\sim 1/a$. This means that the heavy quark fields in the Symanzik action will have no degree of freedom corresponding to “taste.”

In the limit $am_Q \gg 1$, however, it will not be possible to neglect the doublers with either the Fermilab or the NQRCD actions. That is because these actions approach a static lattice theory, which has no intrinsic suppression of spatial doublers: The energy of a static quark is independent of its three-momentum. This would not be a problem if the light quark in

¹ The formalism will also apply when the heavy quark is simulated by extrapolation up in mass from a lighter, conventional, lattice quark.

² For convenience, we use the term “doubler” to describe a heavy-quark state with one or more components of 3-momentum near π/a . Its “rest mass” is defined to be the energy of the state when all 3-momentum components are exactly π/a or 0. Since these states are always suppressed in the cases considered here, we do not need to be concerned that the interpretation of such states as additional heavy *particles* is problematic: With Wilson-like or non-relativistic fermions, their energy *decreases* when one of the large components of momentum changes away from π/a .

a heavy-light bound state were simulated as a Wilson-like fermion. In that case, the light-quark Wilson term would suppress gluon exchanges of momentum π/a between the light and heavy quarks, by forcing the light quark far off shell. But emission or absorption of such a gluon is not suppressed by the staggered light action, and indeed is simply a taste-changing interaction. To keep the analysis manageable, we must assume that the *heavy* quark action suppresses these exchanges. We define Δ_M to be the splitting between (the rest masses of) the physical heavy quark and the closest doublers. To be able to ignore the effect of doubler states in the low energy effective theory, we must have $\Delta_M \gg \Lambda_{QCD}$. However, for the Fermilab or NRQCD action, Δ_M goes like $1/(a^2 m_Q)$ for $am_Q \gg 1$; eventually the condition $\Delta_M \gg \Lambda_{QCD}$ will be violated as m_Q grows at fixed a .

We wish to correct a possible misunderstanding here. When $am_Q \sim 1$, one might worry that doublers cannot be neglected relative to the physical heavy quark since both physical and doubler masses are formally $\mathcal{O}(1/a)$. However, what is important is not (say) the ratio of doubler mass to physical mass, but the *splitting* Δ_M between doubler masses and the physical mass. As long as $\Delta_M \gg \Lambda_{QCD}$, the doubler states can be treated as “integrated out” and their effects can be summarized by higher-dimension operators in the Symanzik/HQET theory [2, 20] involving only the light quarks and the *physical* heavy quark. Systematic errors coming from doublers are thus included in estimates of heavy-quark discretization effects from higher operators. (See further discussion below).

It is instructive to estimate Δ_M in some practical situations. For Wilson-type quarks, the rest (or “pole”) mass m_1 of the physical quark is given at tree level by $am_1 = \ln(1 + am_0)$, where $am_0 = 1/(2\kappa) - 1/(2\kappa_{crit})$, with κ the hopping parameter and κ_{crit} its critical value. For the lowest doubler, with momentum π/a in a single lattice direction, the Wilson term $1 - \cos(ap)$ becomes 2 instead of 0, and the doubler rest mass m_1^D is given by $am_1^D = \ln(3 + am_0)$. For the MILC “coarse” lattices [10], $a^{-1} \approx 1.59$ GeV, $\kappa_{crit} \approx 0.1378$, and κ_b , the hopping parameter of a b quark is ≈ 0.086 [26]. This gives $\Delta_M \equiv m_1^D - m_1 \approx 775$ MeV. Alternatively, using the tadpole improvement [27] parameter $u_0 \equiv 1/(8\kappa_{crit})$ gives $am_0 = 4(\kappa_{crit}/\kappa - 1)$ and $\Delta_M \approx 735$ MeV. These values of Δ_M , which are comparable to other masses dropped in χ PT (*e.g.*, m_ρ), are probably large enough to neglect the effect of doublers in the current chiral analysis. However, it is clear that applying this analysis to any lattices that are significantly coarser than the MILC coarse lattices will be problematic. For example, at $a^{-1} = 1$ GeV, we estimate $\Delta_M \approx 350$ MeV, which is certainly not much larger than Λ_{QCD} . With Δ_M this

low, any effective chiral theory becomes very complicated, and there also would be practical problems in separating the physical state from the doubler states in simulations.

A similar analysis applies in NRQCD. Using a bare mass $aM = 2.8$ for b quarks on the MILC coarse lattices and a stabilization parameter $n = 2$ [6], we find $\Delta_M \approx 930$ MeV, which should be adequate. However, with $a^{-1} = 1$ GeV, $\Delta_M \approx 400$ MeV, again much too low to omit the effects of doublers from the effective chiral theory.

If heavy quark doubler effects may be neglected, the analysis of the Symanzik action is very similar to that in Ref. [28], which considers Ginsparg-Wilson valence quarks coupled to staggered sea quarks. Taste violations at $\mathcal{O}(a^2)$ appear only in four-quark operators composed exclusively of light (staggered) quarks. These terms in the Symanzik action are identical to those in [11, 13]. “Mixed” four-quark operators consisting of the product of a light quark bilinear and a heavy quark bilinear do not break taste symmetry. Physically, operators that violate taste symmetry require momentum π/a gluon exchanges, which we ban from the Symanzik action when we omit doubler states from the low-energy theory. A more rigorous proof that follows from the symmetries of the lattice action can also be constructed. The proof is identical to that presented in Ref. [28], so we do not include it in detail here. The basic idea is that the continuum, physical, heavy quark fields transform trivially under translations by one lattice unit. Staggered bilinears with non-singlet taste, on the other hand, get multiplied by phase factors under single-site translations. Therefore mixed four quark operators cannot be translation singlets unless the light bilinear is a singlet under taste.

The mixed four-quark operators are thus irrelevant from the point of view of our heavy-light chiral theory. Such operators are invariant under the light quark chiral symmetries since they do not break taste (or, trivially, flavor) symmetries and are independent of the light quark masses (otherwise, they would have dimension greater than 6). We therefore can classify the discretization errors caused by such operators as “heavy-quark errors.” These errors can be estimated by the methods of Refs. [2, 20, 21]. (See especially the Appendix of Ref.[21] for estimates of heavy quark discretization errors in a practical case.) In fact, one expects that errors from mixed four-quark operators will be rather smaller than the more familiar ones from heavy-quark bilinear operators, since the former are $\mathcal{O}(\alpha_S^2 a^2)$ for improved staggered (“Asqtad”) light quarks [29]; while the latter are $\mathcal{O}(\alpha_S a)$ or $\mathcal{O}(a^2)$. It is important to keep in mind, however, that such error estimates are based on power counting

and dimensional analysis and must be considered as rough guides only. Precise quantification of the total discretization error will always require simulation at several lattice spacings.

Four-quark operators built entirely out of heavy quarks are also invariant (trivially) under the light quark symmetries. Therefore, the arguments of the previous paragraph apply, with the exception that the discretization errors are now $\mathcal{O}(\alpha_S a^2)$: There is no suppression of the exchange of a single gluon with momentum π/a between the two quarks, unlike the cases where one or both of the quarks has the Asqtad action.

The relevant part of the Symanzik action may now be written as $S^{\text{Sk}} = \int d^4x \mathcal{L}^{\text{Sk}}$, with

$$\mathcal{L}^{\text{Sk}} = \mathcal{L}_4^{\text{Sk}} + a^2 \mathcal{L}_6^{\text{Sk}} + \mathcal{O}(a^4) , \quad (1)$$

where $\mathcal{L}_6^{\text{Sk}}$ comes from light four-quark operators only. $\mathcal{L}_4^{\text{Sk}}$ is the continuum limit of the lattice theory with n staggered light quarks (each with 4 tastes), and a single heavy quark. It has the form

$$\mathcal{L}_4^{\text{Sk}} = \sum_{j=1}^n \bar{q}_j (i \not{D} - m_j) q_j + \bar{Q} (i v \cdot D) Q + \mathcal{L}_{\text{gluons}} . \quad (2)$$

Here v is the heavy quark velocity, and j runs over the light quark flavors. $\mathcal{L}_{\text{Sk},4}$ has a heavy quark $SU(2)$ spin symmetry and, in the limit that $m_j \rightarrow 0$, a chiral $SU(4n)_L \times SU(4n)_R$ symmetry.

The $\mathcal{O}(a^2)$ term $\mathcal{L}_6^{\text{Sk}}$ in Eq. (1) is a sum over all 4-light-quark operators in the two classes $S_6^{FF(A)}$ and $S_6^{FF(B)}$ identified in Ref. [11]:

$$\mathcal{L}_6^{\text{Sk}} = \mathcal{L}_A^{\text{Sk}} + \mathcal{L}_B^{\text{Sk}} . \quad (3)$$

$S_6^{FF(A)}$ consists of ten operators that do not violate continuum rotation symmetry, while the four operators in $S_6^{FF(B)}$ violate this symmetry. For n staggered fields, the operators in each of these classes are written down in Ref. [30]. Every operator is a product of two bilinears, each of which has the same spin and taste. Denoting the five possible spins structures by S , V , T , A , P , and similarly for tastes, the operators are named by the spin $\otimes$ taste of their bilinears:

$$\begin{aligned} \mathcal{L}_A^{\text{Sk}} : \quad & [S \times A], [S \times V], [A \times S], [V \times S], [P \times A], [P \times V], \\ & [A \times P], [V \times P], [T \times V], [T \times A], [V \times T], [A \times T] \end{aligned} \quad (4)$$

$$\mathcal{L}_B^{\text{Sk}} : \quad [T_\mu \times V_\mu], [T_\mu \times A_\mu], [V_\mu \times T_\mu], [A_\mu \times T_\mu] \quad (5)$$

In $\mathcal{L}_A^{\text{Sk}}$ the spin indices and taste indices are contracted separately, for example,

$$[T \times A] \equiv \sum_{\mu < \nu} \sum_{\lambda} \bar{q}_i(\gamma_{\mu\nu} \otimes \xi_{\lambda 5}) q_i \bar{q}_j(\gamma_{\nu\mu} \otimes \xi_{5\lambda}) q_j \quad (6)$$

where $\gamma_{\mu\nu} \equiv (1/2)[\gamma_\mu, \gamma_\nu]$, $\gamma_{\mu 5} \equiv \gamma_\mu \gamma_5$ (similarly for tastes, with $\gamma_\mu \rightarrow \xi_\mu$), and there are implicit sums over the flavor indices i, j . The operators in $\mathcal{L}_B^{\text{Sk}}$ are more complicated: spin and taste indices are summed together, and two terms, with different treatment of the tensor indices, are subtracted to cancel taste- or spin-singlet contributions. For example,

$$[T_\mu \times A_\mu] \equiv \sum_{\mu} \sum_{\mu \neq \nu} \left\{ \bar{q}_i(\gamma_{\mu\nu} \otimes \xi_{\mu 5}) q_i \bar{q}_j(\gamma_{\nu\mu} \otimes \xi_{5\mu}) q_j - \bar{q}_i(\gamma_{\mu\nu 5} \otimes \xi_{\mu 5}) q_i \bar{q}_j(\gamma_{5\nu\mu} \otimes \xi_{5\mu}) q_j \right\} \quad (7)$$

Because the same index (μ) appears four times in each term in $S_6^{FF(B)}$, these operators are not invariant under continuum Euclidean rotations or taste transformations, but only under the lattice symmetry where taste and Euclidean rotations by 90° are performed simultaneously.

In Refs. [11] and [13], the leading order (LO) chiral Lagrangian for light-light pseudoscalars is found for the one-flavor and many-flavor cases, respectively. There the only available Lorentz 4-vector in the chiral theory is the partial derivative operator ∂_μ . To construct a chiral operator with the symmetries of $\mathcal{L}_B^{\text{Sk}}$ operators requires at least two derivatives and two explicit taste indices. Such operators are $\mathcal{O}(p^2 a^2)$, where the factors of lattice spacing a come from the explicit violation of taste symmetry, and the factors of p come from the derivatives. These operators then only occur at next-to-leading-order (NLO) in the chiral Lagrangian, unlike the representatives of $\mathcal{L}_A^{\text{Sk}}$ which are LO. The $\mathcal{L}_B^{\text{Sk}}$ representatives are therefore omitted in Refs. [11, 13], although they are included in Ref [30], which works to NLO.

In the heavy-light case, however, there are two 4-vectors available in the chiral theory whose presence does not raise the order of the operator: the heavy quark 4-velocity v_μ and the light quark gamma matrix γ_μ . Therefore the chiral representatives of the $\mathcal{L}_A^{\text{Sk}}$ operators contribute at the same order as the representatives of the $\mathcal{L}_B^{\text{Sk}}$. For heavy-light decay constants this is NLO — such operators enter into analytic terms in the calculation presented here, but not in one-loop diagrams.

III. THE χ Pt LAGRANGIAN WITH HEAVY-LIGHT MESONS

A. Leading order continuum theory

We first discuss standard χ PT at LO for heavy-light mesons in the continuum [31, 32, 33, 34]. More precisely in the current development, this is the LO chiral representative of the dimension 4 part of the Symanzik Lagrangian, $\mathcal{L}_4^{\text{Sk}}$, Eq. (2). Due to the heavy quark spin symmetry in the static limit, the heavy vector and pseudoscalar mesons are incorporated into the following field, which destroys a heavy-light meson

$$H_a = \frac{1 + \not{v}}{2} [\gamma^\mu B_{\mu a}^* + i\gamma_5 B_a] , \quad (8)$$

where v is the meson's velocity, and a is the combined light quark flavor-taste index. The conjugate field creates a heavy-light meson

$$\overline{H}_a \equiv \gamma_0 H_a^\dagger \gamma_0 = [\gamma^\mu B_{\mu a}^{\dagger*} + i\gamma_5 B_a^\dagger] \frac{1 + \not{v}}{2} , \quad (9)$$

We use B to denote a generic pseudoscalar heavy meson and B^* to denote the corresponding vector meson, but note that the current formalism will also apply to D and D^* mesons, although at a decreased level of accuracy in the $1/m_Q$ expansion.

Under the $SU(2)$ heavy quark spin symmetry, the heavy-light field transforms as

$$\begin{aligned} H &\rightarrow SH , \\ \overline{H} &\rightarrow \overline{H}S^\dagger , \end{aligned} \quad (10)$$

with $S \in SU(2)$, while under the $SU(4n)_L \times SU(4n)_R$ chiral symmetry,

$$\begin{aligned} H &\rightarrow H\mathbb{U}^\dagger , \\ \overline{H} &\rightarrow \mathbb{U}\overline{H} , \end{aligned} \quad (11)$$

with $\mathbb{U} \in SU(4n)$, and we are keeping the flavor-taste index implicit.

We use the term “pion” generically to denote any of the light pseudoscalar mesons. The field describing the pions is $\Sigma = \exp[i\Phi/f]$, where f is the tree-level pion decay constant ($f \sim f_\pi \cong 131$ MeV). In terms involving the heavy-lights, we also need $\sigma \equiv \sqrt{\Sigma} = \exp[i\Phi/2f]$. These fields are singlets under the heavy-quark spin symmetry, while under $SU(4n)_L \times SU(4n)_R$ we have

$$\Sigma \rightarrow L\Sigma R^\dagger , \quad \Sigma^\dagger \rightarrow R\Sigma^\dagger L^\dagger , \quad (12)$$

$$\sigma \rightarrow L\sigma\mathbb{U}^\dagger = \mathbb{U}\sigma R^\dagger , \quad \sigma^\dagger \rightarrow R\sigma^\dagger\mathbb{U}^\dagger = \mathbb{U}\sigma^\dagger L^\dagger , \quad (13)$$

with $L \in SU(4n)_L$ and $R \in SU(4n)_R$. Equation (13) defines $\mathbb{U}$, which is a function of the pion fields and the coordinates, as well as of the global transformations L and R . Chiral operators are formed from combinations of $H, \overline{H}, \sigma, \sigma^\dagger$ and derivatives, as well other matrices such as the light quark mass matrix or taste matrices, so that they transform in the same manner under the combined symmetry group as the underlying operators in the Symanzik action. As usual, this is accomplished by promoting any symmetry-violating factors in the Symanzik Lagrangian to spurions and choosing the transformation properties of the spurions to make the Lagrangian invariant.

For n KS flavors, $\Sigma = \exp(i\Phi/f)$ is a $4n \times 4n$ matrix, and Φ is given by:

$$\Phi = \begin{pmatrix} U & \pi^+ & K^+ & \cdots \\ \pi^- & D & K^0 & \cdots \\ K^- & \bar{K}^0 & S & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad (14)$$

where $U = \sum_{\Xi=1}^{16} U_\Xi T_\Xi$, *etc.*, with the Hermitian taste generators T_Ξ given by

$$T_\Xi = \{\xi_5, i\xi_{\mu 5}, i\xi_{\mu\nu}, \xi_\mu, \xi_I\}. \quad (15)$$

As in Ref. [13] the taste matrices are ξ_μ , with $\xi_{\mu 5} \equiv \xi_\mu \xi_5$. The 4×4 identity matrix is $\xi_I \equiv I$. In this paper, we define $\xi_{\mu\nu} \equiv (1/2)[\xi_\mu, \xi_\nu]$ (rather than simply $\xi_\mu \xi_\nu$) to emphasize that terms with $\mu = \nu$ are never included; in Eq. (15) we take $\mu < \nu$ only.

The component fields of the diagonal (flavor-neutral) elements (U_Ξ, D_Ξ , *etc.*) are real; the other (charged) fields are complex ($\pi_\Xi^\pm, K_\Xi^0$, *etc.*), such that Φ is Hermitian. Here the $n = 3$ portion of Φ is shown explicitly. The mass matrix is the $4n \times 4n$ matrix

$$\mathcal{M} = \begin{pmatrix} m_u I & 0 & 0 & \cdots \\ 0 & m_d I & 0 & \cdots \\ 0 & 0 & m_s I & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad (16)$$

where again, the portion shown is for the $n = 3$ case.

It is convenient to define objects involving the σ field that transform only with $\mathbb{U}$ and $\mathbb{U}^\dagger$. The two possibilities with a single derivative are

$$\mathbb{V}_\mu = \frac{i}{2} [\sigma^\dagger \partial_\mu \sigma + \sigma \partial_\mu \sigma^\dagger], \quad (17)$$

$$\mathbb{A}_\mu = \frac{i}{2} [\sigma^\dagger \partial_\mu \sigma - \sigma \partial_\mu \sigma^\dagger], \quad (18)$$

$\mathbb{V}_\mu$ transforms like a vector field under the $SU(4n)_L \times SU(4n)_R$ chiral symmetry and, when combined with the derivative, can form a covariant derivative acting on the heavy-light field or its conjugate:

$$\begin{aligned} (H\overleftarrow{D}_\mu)_a &= H_b\overleftarrow{D}_\mu^{ba} \equiv \partial_\mu H_a + iH_b\mathbb{V}_\mu^{ba} , \\ (\overrightarrow{D}_\mu\overline{H})_a &= \overrightarrow{D}_\mu^{ab}\overline{H}_b \equiv \partial_\mu\overline{H}_a - i\mathbb{V}_\mu^{ab}\overline{H}_b , \end{aligned} \quad (19)$$

with implicit sums over repeated indices. The covariant derivatives and $\mathbb{A}_\mu$ transform under the chiral symmetry as

$$\begin{aligned} H\overleftarrow{D}_\mu &\rightarrow (H\overleftarrow{D}_\mu)\mathbb{U}^\dagger , \\ \overrightarrow{D}_\mu\overline{H} &\rightarrow \mathbb{U}(\overrightarrow{D}_\mu\overline{H}) , \\ \mathbb{A}_\mu &\rightarrow \mathbb{U}\mathbb{A}_\mu\mathbb{U}^\dagger . \end{aligned} \quad (20)$$

We then write the (Minkowski space) continuum LO chiral Lagrangian as

$$\mathcal{L}_{\text{LO,cont}} = \mathcal{L}_{\text{LO,cont}}^{\text{pion}} + \mathcal{L}_1 , \quad (21)$$

where $\mathcal{L}_{\text{LO,cont}}^{\text{pion}}$ is the standard LO light meson Lagrangian in the continuum, and $\mathcal{L}_1$ is the leading term involving the heavy-lights. As explained in Sec. IIIB, lattice corrections affect the heavy-light terms only at NLO, so there is no need to specify “cont” on $\mathcal{L}_1$. We have

$$\mathcal{L}_{\text{LO,cont}}^{\text{pion}} = \frac{f^2}{8} \text{Tr}(\partial_\mu\Sigma\partial^\mu\Sigma^\dagger) + \frac{1}{4}\mu f^2 \text{Tr}(\mathcal{M}\Sigma + \mathcal{M}\Sigma^\dagger) - \frac{2m_0^2}{3}(U_I + D_I + S_I + \dots)^2 \quad (22)$$

and [34]

$$\mathcal{L}_1 = -i \text{Tr}(\overline{H}Hv\cdot\overleftarrow{D}) + g_\pi \text{Tr}(\overline{H}H\gamma^\mu\gamma_5\mathbb{A}_\mu) . \quad (23)$$

Tr means the complete trace over flavor-taste indices and, where relevant, Dirac indices. Since $\overline{H}$ and H always appear together in the Lagrangian, we treat $\overline{H}H$ as a matrix in flavor-taste space: $(\overline{H}H)_{ab} \equiv \overline{H}_a H_b$. Here and below, the covariant derivative $\overleftarrow{D}$ acts only on the field (in this case H) immediately preceding it; similarly $\overrightarrow{D}$ acts only on the field immediately following it. As in Ref. [13], it is helpful for analyzing the quark flow to leave the anomaly (m_0^2) term explicit in Eq. (22) and work with the diagonal fields $U, D, S, \dots$, rather than the physical ones $\pi_0, \eta, \dots$. At the end of the calculation, we can take $m_0^2 \rightarrow \infty$ and return to the physical basis [35].

Although fundamentally we are interested in the chiral effective theory for a *Euclidean* lattice theory, we find it more convenient to write the Lagrangian in Minkowski space to

make direct contact with the extensive continuum literature. A Wick rotation to define the Euclidean Green's functions is implicit everywhere.

For the heavy-light decay constants, we also need the chiral representative of the axial heavy-light current. Alternatively, one can work with the left-handed current, which has the advantage that it transforms more simply under chiral transformations and in addition gives corresponding information about the vector current, useful for semileptonic form factors. The left-handed current that destroys a heavy-light meson of flavor-taste b is $j^{\mu,b}$, which at LO takes the form [34]

$$j_{\text{LO}}^{\mu,b} = \frac{\kappa}{2} \text{tr}_D(\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} \quad (24)$$

where $\lambda^{(b)}$ is a constant vector that fixes the flavor-taste: $(\lambda^{(b)})_a = \delta_{ab}$, and tr_D is a trace on Dirac indices only. In QCD, the decay constant f_{B_a} is defined by the matrix element

$$\langle 0 | j^{\mu,b} | B_a(v) \rangle = i f_{B_a} m_{B_a} v^\mu \delta_{ab} , \quad (25)$$

where relativistic normalization of the state $|B_a(v)\rangle$ is assumed. At LO in the heavy-light chiral theory, $j_{\text{LO}}^{\mu,b} = i\kappa v^\mu B_b$, which gives $f_{B_a}^{\text{LO}} = \kappa/\sqrt{m_{B_a}}$. Recall that the factor $\sqrt{m_{B_a}}$ arises from the differences in normalizations between relativistic and non-relativistic states.

B. Power counting

Before considering discretization errors and higher order corrections to Eqs. (21) and (24), we discuss the power counting assumed in this paper. The staggered chiral Lagrangian with only light mesons is a joint expansion in the light quark mass m_q and the lattice spacing a^2 . In that case, m_q and a^2 are taken to be of the same order in the expansion [11, 13]. Since $m_\pi^2 \propto m_q$ and the momentum of external light mesons, p_π , is assumed to be of order m_π , we have $p_\pi^2 \sim m_\pi^2 \sim m_q \sim a^2$ in the power counting.

Upon including B mesons, we have an additional expansions in the inverse of the heavy quark mass m_Q , and in the B meson's residual momentum, k , which we take to be of the same order as p_π . We work to leading order in $1/m_Q$ only.

The continuum LO chiral Lagrangian, Eq. (21), is therefore $\mathcal{O}(k \sim \sqrt{m_q})$ in the heavy-meson fields and $\mathcal{O}(m_q, a^2)$ in the light meson fields. In each case these are the lowest order terms allowed by the continuum symmetries.³ One-loop diagrams with vertices from

³ The $\mathcal{O}(k^0)$ heavy quark mass term is absorbed by measuring heavy-light energies relative to the mass of

Eqs. (21) and (24) and propagators from Eq. (21) will give $\mathcal{O}(m_\pi^2)$ corrections to physical quantities such as f_{B_a} or semileptonic form factors. Since taste-violating terms of $\mathcal{O}(a^2)$ in the pion Lagrangian of S χ PT are the same order as those already included in Eq. (22), we will need to include such terms in order to calculate the one-loop corrections.

Terms that are higher order than those in Eqs. (21) and (24) are irrelevant in the one-loop diagrams. They can, however, make analytic (tree-level) contributions at this (or, in principle, lower) order. In particular, for f_{B_a} we will need to consider $\mathcal{O}(\sqrt{m_q})$ and $\mathcal{O}(m_q)$ corrections to the existing heavy-light terms in Eqs. (21) and (24). Such terms in the heavy-light Lagrangian, Eq. (23) are of order $m_q \sim a^2$ and $km_q \sim ka^2$; those in the current, Eq. (24), are of order $m_q \sim a^2$. These corrections are generated by inserting, into heavy-light chiral operators, spurions representing the light-quark mass term in $\mathcal{L}_4^{\text{Sk}}$ [Eq. (2)] and the $\mathcal{O}(a^2)$ taste-violating terms, $\mathcal{L}_6^{\text{Sk}}$ [Eq. (3)]. There are also invariant higher order operators that are present even in the continuum and chiral limits. These are generated by inserting additional derivative operators, *i.e.*, additional powers of k . As explained in Sec. III D, such terms do not affect the decay constants to the order we are working, and we do not attempt to catalog them completely. We also do not need to consider here higher order [$\mathcal{O}(m_q^2 \sim m_q a^2 \sim a^4)$] corrections to the pion Lagrangian. Such corrections cannot give analytic contributions to heavy-light decay constants. They are, however, relevant to purely light-light physics, as well as to semileptonic form factors of heavy-light mesons, which involve a light-light particle. All such terms have been found in Ref. [30].

Summarizing the discussion of this subsection, we write our complete chiral Lagrangian as

$$\mathcal{L} = \mathcal{L}_{\text{LO}} + \mathcal{L}_{\text{NLO}} , \quad (26)$$

$$\mathcal{L}_{\text{LO}} = \mathcal{L}_{\text{LO}}^{\text{pions}} + \mathcal{L}_1 = \mathcal{L}_{\text{LO,cont}}^{\text{pions}} + \mathcal{L}_{\text{LO},a^2}^{\text{pions}} + \mathcal{L}_1 , \quad (27)$$

$$\mathcal{L}_{\text{NLO}} = \mathcal{L}_2 + \mathcal{L}_3 \quad (28)$$

$$\mathcal{L}_2 = \mathcal{L}_{2,k} + \mathcal{L}_{2,m} + \mathcal{L}_{2,a^2}^A + \mathcal{L}_{2,a^2}^B , \quad (29)$$

$$\mathcal{L}_3 = \mathcal{L}_{3,k} + \mathcal{L}_{3,m} + \mathcal{L}_{3,a^2}^A + \mathcal{L}_{3,a^2}^B . \quad (30)$$

Here, $\mathcal{L}_1$, $\mathcal{L}_2$, and $\mathcal{L}_3$ involve the heavy-light fields and are of order k , $k^2 \sim m_q \sim a^2$, and $k^3 \sim km_q \sim ka^2$, respectively. The subscripts k , m , or a^2 indicate terms that involve

the heavy meson.

derivatives only, mass spurions, or taste-violating spurions. The superscripts A and B stand for the type of taste-violating terms, Eqs. (4) and (5).

Similarly we have for the left-handed current:

$$j^{\mu,b} = j_{\text{LO}}^{\mu,b} + j_{\text{NLO}}^{\mu,b} , \quad (31)$$

$$j_{\text{NLO}}^{\mu,b} = j_1^{\mu,b} + j_2^{\mu,b} \quad (32)$$

$$j_1^{\mu,b} = j_{1,k}^{\mu,b} \quad (33)$$

$$j_2^{\mu,b} = j_{2,k}^{\mu,b} + j_{2,m}^{\mu,b} + j_{2,a^2,A}^{\mu,b} + j_{2,a^2,B}^{\mu,b} . \quad (34)$$

Note that insertions of the mass and a^2 spurions are always at least two orders higher in $k \sim \sqrt{m_q}$, so do not appear in $j_1^{\mu,b}$.

The continuum LO terms $\mathcal{L}_1$, $\mathcal{L}_{\text{LO},\text{cont}}^{\text{pions}}$, and $j_{\text{LO}}^{\mu,b}$ are given in Eqs. (22), (23) and (24). The remaining contributions, Lagrangian terms $\mathcal{L}_{\text{LO},a^2}^{\text{pions}}$, $\mathcal{L}_2$, and $\mathcal{L}_3$, and current terms $j_1^{\mu,b}$ and $j_2^{\mu,b}$, are discussed in the following subsections.

C. LO discretization effects

The LO finite lattice spacing correction, $\mathcal{L}_{\text{LO},a^2}^{\text{pion}}$, to the pion Lagrangian is known from previous work [11, 13]. We have

$$\begin{aligned} \mathcal{L}_{\text{LO},a^2}^{\text{pion}} &= -a^2 \mathcal{V}_\Sigma \\ \mathcal{V}_\Sigma &= C_1 \text{Tr}(\xi_5^{(n)} \Sigma \xi_5^{(n)} \Sigma^\dagger) + C_3 \frac{1}{2} \sum_\nu [\text{Tr}(\xi_\nu^{(n)} \Sigma \xi_\nu^{(n)} \Sigma) + h.c.] \\ &\quad + C_4 \frac{1}{2} \sum_\nu [\text{Tr}(\xi_{\nu 5}^{(n)} \Sigma \xi_{5\nu}^{(n)} \Sigma) + h.c.] + C_6 \sum_{\mu < \nu} \text{Tr}(\xi_{\mu\nu}^{(n)} \Sigma \xi_{\nu\mu}^{(n)} \Sigma^\dagger) \\ &\quad + C_{2V} \frac{1}{4} \sum_\nu [\text{Tr}(\xi_\nu^{(n)} \Sigma) \text{Tr}(\xi_\nu^{(n)} \Sigma) + h.c.] + C_{2A} \frac{1}{4} \sum_\nu [\text{Tr}(\xi_{\nu 5}^{(n)} \Sigma) \text{Tr}(\xi_{5\nu}^{(n)} \Sigma) + h.c.] \\ &\quad + C_{5V} \frac{1}{2} \sum_\nu \text{Tr}(\xi_\nu^{(n)} \Sigma) \text{Tr}(\xi_\nu^{(n)} \Sigma^\dagger) + C_{5A} \frac{1}{2} \sum_\nu \text{Tr}(\xi_{\nu 5}^{(n)} \Sigma) \text{Tr}(\xi_{5\nu}^{(n)} \Sigma^\dagger) . \end{aligned} \quad (35)$$

The explicit $4n \times 4n$ matrices $\xi_\mu^{(n)}$ in Eq. (35) are defined by

$$(\xi_\nu^{(n)})_{ij} = \xi_\nu \delta_{ij} , \quad (36)$$

with i and j the $SU(n)$ (light quark) flavor indices, and ξ_ν the 4×4 taste matrix, as in Eq. (15). The matrices $\xi_{\mu\nu}^{(n)}$ and $\xi_{\nu 5}^{(n)}$ are defined similarly.

There are no a^2 corrections at LO to $\mathcal{L}_1$, Eq. (23), or to the left-handed current, Eq. (24).

D. Invariant NLO corrections; $\mathcal{T}$ symmetry

Even in the chiral and continuum limit, there are invariant terms that can be constructed by adding more derivatives to the lowest-order heavy-light Lagrangian $\mathcal{L}_1$, or current $j_{\text{LO}}^{\mu,b}$. To our knowledge, a complete catalog of such terms through $\mathcal{O}(k^3)$ does not appear in the continuum literature. These higher derivative terms are in fact irrelevant for the heavy-light decay constant to the order we are working: Derivatives acting on the light-light fields Σ or σ do not contribute to tree diagrams because there are no external pions in the matrix element; while terms with a derivative acting on the heavy-light field in the current, or more than one derivative acting on a heavy-light field in the Lagrangian, give a vanishing contribution on shell ($k = 0$).⁴

For semileptonic form factors, *e.g.*, those for $B \rightarrow \pi$, higher derivative terms do produce analytic corrections at this order. However, the functional form that such contributions can take is rather limited: In the chiral limit, one can only get corrections to form factors proportional to $v \cdot p$ or $(v \cdot p)^2$, where p is the pion 4-momentum. Away from the chiral limit, there will be additional corrections proportional to p^2 , but these merely duplicate, on-shell, the effects of mass or a^2 spurions.

We thus simply list here a few representative operators that appear in $\mathcal{L}_{2,k}$, $\mathcal{L}_{3,k}$, $j_{1,k}^{\mu,b}$, and $j_{2,k}^{\mu,b}$:

$$\mathcal{L}_{2,k} = \frac{i\epsilon_1}{\Lambda_\chi} \text{Tr} \left((v \cdot \vec{D} \overline{H} H - \overline{H} H v \cdot \vec{D}) \gamma_\mu \gamma_5 \mathbb{A}^\mu \right) + \frac{\epsilon_2}{\Lambda_\chi} \text{Tr} \left(\overline{H} (v \cdot \vec{D})^2 H \right) + \dots \quad (37)$$

$$\mathcal{L}_{3,k} = \frac{\epsilon_3}{\Lambda_\chi^2} \text{Tr} \left(\overline{H} H \gamma_\mu \gamma_5 (v \cdot \vec{D})^2 \mathbb{A}^\mu \right) + \frac{\epsilon_4}{\Lambda_\chi^2} \text{Tr} \left(\overline{H} H \vec{D} \gamma_5 v \cdot \vec{D} v \cdot \mathbb{A} \right) + \dots \quad (38)$$

$$j_{1,k}^{\mu,b} = \frac{i\kappa_1}{\Lambda_\chi} \text{tr}_D \left(\gamma^\mu (1 - \gamma_5) v \cdot \vec{D} H \right) \sigma^\dagger \lambda^{(b)} + \frac{\kappa_2}{\Lambda_\chi} \text{tr}_D \left(\gamma^\mu (1 - \gamma_5) H \right) v \cdot \mathbb{A} \sigma^\dagger \lambda^{(b)} + \dots \quad (39)$$

$$j_{2,k}^{\mu,b} = \frac{\kappa_3}{\Lambda_\chi^2} \text{tr}_D \left(\gamma^\mu (1 - \gamma_5) (v \cdot \vec{D})^2 H \right) \sigma^\dagger \lambda^{(b)} + \frac{i\kappa_4}{\Lambda_\chi^2} \text{tr}_D \left(\gamma^\mu (1 - \gamma_5) H \right) v \cdot \vec{D} v \cdot \mathbb{A} \sigma^\dagger \lambda^{(b)} + \dots \quad (40)$$

where the constants ϵ_i, κ_j are taken to be real and dimensionless, Λ_χ is the chiral scale, and

$$\vec{D}_\nu \mathbb{A}_\mu \equiv \partial_\nu \mathbb{A}_\mu - i[\mathbb{V}_\nu, \mathbb{A}_\mu] . \quad (41)$$

⁴ Terms with a single derivative in the Lagrangian do contribute on shell through wave-function renormalization.

The use of time reversal ($\mathcal{T}$) symmetry and the requirement of Hermiticity for the Lagrangian can be used to eliminate several other candidate operators from Eqs. (37) through (40). Because we will also need to use the consequences of $\mathcal{T}$ invariance extensively in Secs. III E to IV B, we briefly review how it acts on relevant objects, following Ref. [36]. Let $\tilde{p}$ be the parity reflection of a 4-vector p ; in other words $\tilde{p}^\mu = p_\mu$. Then time reversal acts as follows

$$\begin{aligned}
x &\rightarrow -\tilde{x} \\
v &\rightarrow \tilde{v} \\
\Phi(x) &\rightarrow -\Phi(-\tilde{x}) \\
\Sigma(x) &\rightarrow \Sigma(-\tilde{x}) \\
\sigma(x) &\rightarrow \sigma(-\tilde{x}) \\
\mathbb{A}^\mu(x) &\rightarrow \mathbb{A}_\mu(-\tilde{x}) \\
\mathbb{V}^\mu(x) &\rightarrow \mathbb{V}_\mu(-\tilde{x}) \\
\vec{D}^\mu &\rightarrow -\vec{D}_\mu \\
H^v(x) &\rightarrow T H^{\tilde{v}}(-\tilde{x}) T^{-1} \\
\overline{H}^v(x) &\rightarrow T \overline{H}^{\tilde{v}}(-\tilde{x}) T^{-1}
\end{aligned} \tag{42}$$

where the Dirac matrix T has the property $T\gamma^\mu T^{-1} = \gamma_\mu^*$, we show the v dependence of H explicitly, and we have used the anti-unitary nature of $\mathcal{T}$.

Requiring $\mathcal{T}$ invariance, we can for example eliminate contributions to the Lagrangian such as

$$\text{Tr} \left(\overline{H} H \gamma_\mu \gamma_5 v \cdot \vec{D} \mathbb{A}^\mu \right) . \tag{43}$$

Adding a factor of i to make this $\mathcal{T}$ -even, as was done in Ref. [36], does not in fact save this term because it is then anti-Hermitian.

E. Quark mass corrections at NLO

Here we discuss operators induced by single insertions of the light quark mass spurions in $\mathcal{L}_4^{Sk}$. These produce corrections to the Lagrangian, namely $\mathcal{L}_{2,m}$ and $\mathcal{L}_{3,m}$, and corrections to the current, $j_{2,m}^{\mu,b}$. Of course, such terms are also present at NLO in the continuum, and the result is more or less standard [36, 37]. However, since many of the contributions to

the discretization corrections ($\mathcal{L}_{2,a^2}^A, \mathcal{L}_{2,a^2}^B, \mathcal{L}_{3,a^2}^A, \mathcal{L}_{3,a^2}^B, j_{2,a^2,A}^{\mu,b}$, and $j_{2,a^2,B}^{\mu,b}$) follow the same pattern (see Secs. IV A and IV B), it is worthwhile to examine the mass corrections carefully first. In addition, we find some modifications to the terms written down in Refs. [36, 37].

We start by writing down combinations of the light quark mass spurions that transform only with $\mathbb{U}$ or $\mathbb{U}^\dagger$. These are

$$\mathcal{M}^\pm = \frac{1}{2} (\sigma \mathcal{M} \sigma \pm \sigma^\dagger \mathcal{M} \sigma^\dagger) , \quad (44)$$

which transform as $\mathcal{M}^\pm \rightarrow \mathbb{U} \mathcal{M}^\pm \mathbb{U}^\dagger$ under the chiral symmetry. Under parity, $\mathcal{M}^\pm \rightarrow \pm \mathcal{M}^\pm$, since $\Phi \rightarrow -\Phi$ and $\sigma \rightarrow \sigma^\dagger$; while both $\mathcal{M}^+$ and $\mathcal{M}^-$ are even under time reversal. Because $\text{tr}_D(\overline{H} H \gamma_5) = 0$, the only parity and chiral invariants we can construct with no derivatives are $\text{Tr}(\overline{H} H \mathcal{M}^+)$ and $\text{Tr}(\overline{H} H) \text{Tr}(\mathcal{M}^+)$. We thus have

$$\mathcal{L}_{2,m} = 2\lambda_1 \text{Tr}(\overline{H} H \mathcal{M}^+) + 2\lambda'_1 \text{Tr}(\overline{H} H) \text{Tr}(\mathcal{M}^+) . \quad (45)$$

For $\mathcal{L}_{3,m}$, we start with the lists of terms given in Eq. (16) of Ref. [36] and Eq. (7) of Ref. [37]. Note, however, that a term like $\text{Tr}(\overline{H} H \gamma_\mu \gamma_5 \mathbb{A}^\mu \mathcal{M}^+)$ is not Hermitian, because the Hermitian conjugate will flip the order of $\mathcal{M}^+$ and $\mathbb{A}^\mu$. To make it Hermitian we can write it in one of two ways

$$\begin{aligned} & \text{Tr}(\overline{H} H \gamma_\mu \gamma_5 \{\mathbb{A}^\mu, \mathcal{M}^+\}) , \\ & i \text{Tr}(\overline{H} H \gamma_\mu \gamma_5 [\mathbb{A}^\mu, \mathcal{M}^+]) . \end{aligned}$$

The second term is not invariant under time-reversal and can be dropped. For corresponding terms with $\mathcal{M}^-$, it is the anticommutator that requires a factor of i , since $\mathcal{M}^-$ is anti-Hermitian. Then time-reversal invariance requires that we build the traces using the commutator $[\mathbb{A}^\mu, \mathcal{M}^-]$ only. Terms with separately traced operators, such as $\text{Tr}(\overline{H} H \gamma_\mu \gamma_5) \text{Tr}(\mathbb{A}^\mu \mathcal{M}^+)$, do not need to be rewritten with commutators or anticommutators since they are Hermitian already.

Terms involving a derivative acting on the heavy-light fields also require some thought. Hermitian combinations are

$$\begin{aligned} & i \left(\overrightarrow{D}_\mu \overline{H} H - \overline{H} H \overleftarrow{D}_\mu \right) , \\ & \overrightarrow{D}_\mu \overline{H} H + \overline{H} H \overleftarrow{D}_\mu . \end{aligned}$$

However, we will not need to include the second combination, which is a total derivative. When combined a factor of $\mathcal{M}^\pm$, integration by parts can be used to put the derivative on

the σ or $\sigma^\dagger$ fields. Such terms can then be rewritten in terms of $\mathbb{A}$, using $i\vec{D}^\mu\sigma^\dagger = -\mathbb{A}^\mu\sigma^\dagger$ or $i\vec{D}^\mu\sigma = \mathbb{A}^\mu\sigma$ [36].

Putting together the above discussion, we have

$$\begin{aligned}\mathcal{L}_{3,m} = & ik_1 \text{Tr} \left(\overline{H} H v \cdot \overleftarrow{D} \mathcal{M}^+ - v \cdot \overrightarrow{D} \overline{H} H \mathcal{M}^+ \right) \\ & + ik_2 \text{Tr} \left(\overline{H} H v \cdot \overleftarrow{D} - v \cdot \overrightarrow{D} \overline{H} H \right) \text{Tr}(\mathcal{M}^+) \\ & + k_3 \text{Tr} \left(\overline{H} H \gamma_\mu \gamma_5 \{ \mathbb{A}^\mu, \mathcal{M}^+ \} \right) + k_4 \text{Tr} \left(\overline{H} H \gamma_\mu \gamma_5 \mathbb{A}^\mu \right) \text{Tr}(\mathcal{M}^+) \\ & + k_5 \text{Tr} \left(\overline{H} H \gamma_\mu \gamma_5 \right) \text{Tr} \left(\mathbb{A}^\mu \mathcal{M}^+ \right) + k_6 \text{Tr} \left(\overline{H} H \gamma_\mu [\mathbb{A}^\mu, \mathcal{M}^-] \right) .\end{aligned}\quad (46)$$

Note that $\overline{H}$ and H must be next to each other in these traces by heavy quark spin symmetry. The term that involves $\mathcal{M}^-$ will not contribute to the decay constants or to the form factors for semileptonic decays, but we include it here to complete the list of mass-dependent terms.

For the current, we have:

$$\begin{aligned}j_{2,m}^{\mu,b} = & \rho_1 \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \mathcal{M}^+ \sigma^\dagger \lambda^{(b)} + \rho_2 \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} \text{Tr}(\mathcal{M}^+) \\ & + \rho_3 \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \mathcal{M}^- \sigma^\dagger \lambda^{(b)} + \rho_4 \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} \text{Tr}(\mathcal{M}^-) .\end{aligned}\quad (47)$$

It is not hard to argue that the heavy quark spin symmetry forces the factor $\gamma^\mu(1 - \gamma_5)$ to appear next to the field H ; it could not, for example, be replaced with $v^\mu(1 - \gamma_5)$. To show this, one can replace $\gamma^\mu(1 - \gamma_5)$ in the QCD current by a spurion, and give it appropriate transformation properties under heavy quark spin symmetry and Lorentz transformations to make the current invariant. A similar argument will apply even for the more complicated terms in the current described below.

IV. TASTE BREAKING WITH HEAVY-LIGHT MESONS

The final needed contributions to the chiral Lagrangian and current are $\mathcal{O}(a^2)$ terms involving heavy-light fields: $\mathcal{L}_{2,a^2}^A$, $\mathcal{L}_{2,a^2}^B$, $\mathcal{L}_{3,a^2}^A$, $\mathcal{L}_{3,a^2}^B$ [Eqs. (29) and (30)] and $j_{2,a^2,A}^{\mu,b}$, $j_{2,a^2,B}^{\mu,b}$ [Eq. (34)]. These are constructed by starting with the dimension-6 terms in the Symanzik Lagrangian, $\mathcal{L}_A^{\text{Sk}}$ and $\mathcal{L}_B^{\text{Sk}}$ [Eqs. (4) and (5)]. We promote the two explicit taste matrices in each of these operators to spurion fields, which are assigned transformation laws such that the operators are invariant under the chiral $SU(4n)_L \times SU(4n)_R$ symmetry.

The heavy quark spin symmetry requires that we only have the combination $\overline{H}H$ in the

Lagrangian, and $\gamma^\mu(1 - \gamma_5)H$ in the current. Under the chiral symmetry, these transform as

$$\begin{aligned}\overline{H}H &\rightarrow \mathbb{U}\overline{H}H\mathbb{U}^\dagger \\ \gamma^\mu(1 - \gamma_5)H &\rightarrow \gamma^\mu(1 - \gamma_5)H\mathbb{U}^\dagger.\end{aligned}$$

When doing the spurion analysis, we therefore need to combine the taste matrices (treated as spurion fields), σ , and $\sigma^\dagger$ into operators $\mathcal{O}_i$ that obey:

$$\mathcal{O}_i \rightarrow \mathbb{U}\mathcal{O}_i\mathbb{U}^\dagger. \quad (48)$$

We then construct chiral (and heavy-quark spin) invariants using these spurions and the heavy-light fields. At $\mathcal{O}(a^2)$, exactly two taste spurions will appear in each term in the Lagrangian and the current. In $\mathcal{L}_{3,a^2}^A$ and $\mathcal{L}_{3,a^2}^B$, which are $\mathcal{O}(ka^2)$, a derivative term will also be needed. Either D_μ or $\mathbb{A}_\mu$ can be used.

A large class of contributions to $\mathcal{L}_{2,a^2}^A$, $\mathcal{L}_{3,a^2}^A$, and $j_{2,a^2,A}^{\mu,b}$ can be obtained by first combining both taste spurions into a single operator that transforms like Eq. (48). As described in Sec. IV A, all such operators may be easily found from known results using a trick. These operators are the same order in our chiral power counting and have the same transformation properties as the quark mass terms, $\mathcal{M}^\pm$. Thus, we can simply replace the $\mathcal{M}^\pm$ in Eqs. (45), (46) and (47) with such two-taste-spurion operators to find all terms in $\mathcal{L}_{2,a^2}^A$, $\mathcal{L}_{3,a^2}^A$, and $j_{2,a^2,A}^{\mu,b}$ in which the two spurions combine together into a single operator. It is then not hard to catalog the remaining contributions, which are ones in which the two spurions are interspersed between $\overline{H}H$ and $\mathbb{A}$ in a single trace, or are separately traced with $\overline{H}H$ or $\mathbb{A}$.

Determining the chiral representatives of $\mathcal{L}_B^{\text{Sk}}$ works similarly. However each chiral invariant constructed from the spurions will generate several terms in $\mathcal{L}_{2,a^2}^B$, $\mathcal{L}_{3,a^2}^B$, and $j_{2,a^2,B}^{\mu,b}$, due to various ways of introducing the appropriate Lorentz indices using v_μ , γ_μ , D_μ , or $\mathbb{A}_\mu$.

A. Discretization errors at NLO: Operators from $\mathcal{L}_A^{\text{Sk}}$

To find combinations of two taste spurions from $\mathcal{L}_A^{\text{Sk}}$ that transform like Eq. (48), we can start with the invariant light-meson operators that result from the spurion analysis. These are the eight operators given in Eq. (35) [11, 13]. From these operators, we can generate all ones that transform like Eq. (48) by simply replacing a factor of Σ or $\Sigma^\dagger$ with, respectively, σ^2 or $(\sigma^\dagger)^2$, permuting the trace so there is a factor of σ or $\sigma^\dagger$ at both beginning and end of

the term, and removing the trace. For example,

$$\text{Tr}(\xi_5^{(n)} \Sigma^\dagger \xi_5^{(n)} \Sigma) \rightarrow \sigma \xi_5^{(n)} \Sigma^\dagger \xi_5^{(n)} \sigma = \sigma \xi_5^{(n)} \sigma^\dagger \sigma^\dagger \xi_5^{(n)} \sigma , \quad (49)$$

where the final form shows explicitly how the result is composed of the produce of two taste spurions, each transforming like Eq. (48). We call this process “opening up” a Σ (or $\Sigma^\dagger$). It automatically creates an operator that has the desired transformation property when the spurions transform as required. Each operator in Eq. (35) thereby generates two operators, depending on whether one opens up a Σ or a $\Sigma^\dagger$. As was done for the mass term in Eq. (44), the two operators can be joined to make Hermitian or anti-Hermitian combinations, or, equivalently, even or odd parity combinations. This gives us the following operators that transform under chiral rotations by Eq. (48):

$$\begin{aligned} \mathcal{O}_1^{A,\pm} &= (\sigma \xi_5^{(n)} \Sigma^\dagger \xi_5^{(n)} \sigma \pm p.c.) \\ \mathcal{O}_2^{A,\pm} &= \sum_\nu [(\sigma \xi_\nu^{(n)} \sigma) \text{Tr}(\xi_\nu^{(n)} \Sigma) \pm p.c.] \\ \mathcal{O}_3^{A,\pm} &= \sum_\nu (\sigma \xi_\nu^{(n)} \Sigma \xi_\nu^{(n)} \sigma \pm p.c.) \\ \mathcal{O}_4^{A,\pm} &= \sum_\nu (\sigma \xi_{\nu 5}^{(n)} \Sigma \xi_{5\nu}^{(n)} \sigma \pm p.c.) \\ \mathcal{O}_5^{A,\pm} &= \sum_\nu [(\sigma \xi_\nu^{(n)} \sigma) \text{Tr}(\xi_\nu^{(n)} \Sigma^\dagger) \pm p.c.] \\ \mathcal{O}_6^{A,\pm} &= \sum_\mu \sum_{\nu \neq \mu} (\sigma \xi_{\mu\nu}^{(n)} \Sigma^\dagger \xi_{\nu\mu}^{(n)} \sigma \pm p.c.) \\ \mathcal{O}_7^{A,\pm} &= \sum_\nu [(\sigma \xi_{\nu 5}^{(n)} \sigma) \text{Tr}(\xi_{5\nu}^{(n)} \Sigma) \pm p.c.] \\ \mathcal{O}_8^{A,\pm} &= \sum_\nu [(\sigma \xi_{\nu 5}^{(n)} \sigma) \text{Tr}(\xi_{5\nu}^{(n)} \Sigma^\dagger) \pm p.c.] , \end{aligned} \quad (50)$$

where $p.c.$ stands for parity conjugate: for example $\sigma_{p.c.} = \sigma^\dagger$. We have followed the numbering scheme in Eq. (35), with the exceptions that the operators multiplied by C_{2V} and C_{5V} in Eq. (35) become $\mathcal{O}_2^{A,\pm}$ and $\mathcal{O}_5^{A,\pm}$, while those multiplied by C_{2A} and C_{5A} become $\mathcal{O}_7^{A,\pm}$ and $\mathcal{O}_8^{A,\pm}$. We have also not bothered to keep the conventional factors of 1/2 and 1/4 from Eq. (35) in our definitions here. Operators in Eq. (50) with no traces (coming from ones in Eq. (35) with one trace) correspond to multiplying two taste spurions transforming like Eq. (48); operators with a trace (coming from ones in Eq. (35) with two traces) correspond to tracing one taste spurion and then multiplying by the other.

Making the replacement $\mathcal{M}^\pm \rightarrow \mathcal{O}_k^{A,\pm}$ in Eq. (45), we then get

$$\mathcal{L}_{2,a^2}^A = a^2 \sum_{k=1}^8 \left\{ K_{1,k}^A \text{Tr} \left(\overline{H} H \mathcal{O}_k^{A,+} \right) + K_{2,k}^A \text{Tr} \left(\overline{H} H \right) \text{Tr} \left(\mathcal{O}_k^{A,+} \right) \right\} \quad (51)$$

$\mathcal{L}_{3,a^2}^A$ is more complicated, since we can use one factor of $\mathbb{A}^\mu$. Because $\mathbb{A}^\mu$ transforms like $\overline{H} H$, there is now an additional possibility of interspersing single spurions between $\mathbb{A}^\mu$ and $\overline{H} H$. A complete spurion analysis is therefore required for such terms. We can follow Appendix B of Ref. [11], with the simple modifications [13] that we keep only “odd-odd” operators and make the spurions diagonal in flavor, as in Eq. (36). We then prepend and append appropriate factors of σ and $\sigma^\dagger$ to each spurion to make it into an operator that transforms like Eq. (48). The result of this analysis is that a given chiral operator of this kind in $\mathcal{L}_{3,a^2}^A$ will involve a pair of spurion operators, which we call Q_k^A and $\tilde{Q}_k^A$ ($k = 1, \dots, 8$):

$$\begin{aligned} [V \times P], [A \times P] &\rightarrow Q_1^A = \sigma \xi_5^{(n)} \sigma^\dagger, & \tilde{Q}_1^A &\equiv (Q_1^A)_{p.c.} = \sigma^\dagger \xi_5^{(n)} \sigma \\ [V \times P], [A \times P] &\rightarrow Q_2^A = \sigma \xi_5^{(n)} \sigma^\dagger, & \tilde{Q}_2^A &\equiv Q_2^A \\ [S \times V], [P \times V], [T \times V] &\rightarrow Q_3^A = \sigma \xi_\nu^{(n)} \sigma, & \tilde{Q}_3^A &\equiv Q_3^A \\ [S \times A], [P \times A], [T \times A] &\rightarrow Q_4^A = i \sigma \xi_{\nu 5}^{(n)} \sigma, & \tilde{Q}_4^A &\equiv Q_4^A \\ [S \times V], [P \times V] &\rightarrow Q_5^A = \sigma \xi_\nu^{(n)} \sigma, & \tilde{Q}_5^A &\equiv (Q_5^A)_{p.c.} = \sigma^\dagger \xi_\nu^{(n)} \sigma^\dagger \\ [V \times T], [A \times T] &\rightarrow Q_6^A = i \sigma \xi_{\lambda \nu}^{(n)} \sigma^\dagger, & \tilde{Q}_6^A &\equiv (Q_6^A)_{p.c.} = -i \sigma^\dagger \xi_{\nu \lambda}^{(n)} \sigma \\ [V \times T], [A \times T] &\rightarrow Q_7^A = i \sigma \xi_{\lambda \nu}^{(n)} \sigma^\dagger, & \tilde{Q}_7^A &\equiv Q_7^A \\ [S \times A], [P \times A] &\rightarrow Q_8^A = i \sigma \xi_{\nu 5}^{(n)} \sigma, & \tilde{Q}_8^A &\equiv (Q_8^A)_{p.c.} = -i \sigma^\dagger \xi_{5 \nu}^{(n)} \sigma^\dagger, \end{aligned} \quad (52)$$

where we show the terms in $\mathcal{L}_A^{\text{Sk}}$ [Eq. (4)] that generate the given spurion operators. The terms $[V \times S]$ and $[A \times S]$ in the Symanzik action do not appear in this list since the corresponding spurion operators, such as $\sigma \xi_I^{(n)} \sigma^\dagger$, are trivial. We have tried to make the numbering system in Eq. (52) correspond to that in Eqs. (35) and (50) as much as possible. Note that $Q_1^A = Q_2^A$ but $\tilde{Q}_1^A \neq \tilde{Q}_2^A$, and similarly for $Q_3^A = Q_5^A$, $Q_4^A = Q_8^A$, and $Q_6^A = Q_7^A$. Operators obtained by parity conjugating both Q_k^A and $\tilde{Q}_k^A$ are not included since they will appear automatically when we demand parity invariance of the chiral Lagrangian.

We can now construct $\mathcal{L}_{3,a^2}^A$ by first replacing $\mathcal{M}^\pm \rightarrow \mathcal{O}_k^{A,\pm}$ in Eq. (46), and then adding

on new operators that intersperse Q_k^A and $\tilde{Q}_k^A$ between $\overline{H}H$ and $\mathbb{A}^\mu$. The result is:

$$\begin{aligned}
\mathcal{L}_{3,a^2}^A = & a^2 \sum_{k=1}^8 \left\{ i c_{1,k}^A \text{Tr} \left(\overline{H}H v \cdot \overleftarrow{D} \mathcal{O}_k^{A,+} - v \cdot \overrightarrow{D} \overline{H}H \mathcal{O}_k^{A,+} \right) \right. \\
& + i c_{2,k}^A \text{Tr} \left(\overline{H}H v \cdot \overleftarrow{D} - v \cdot \overrightarrow{D} \overline{H}H \right) \text{Tr}(\mathcal{O}_k^{A,+}) \\
& + c_{3,k}^A \text{Tr} \left(\overline{H}H \gamma_\mu \gamma_5 \{ \mathbb{A}^\mu, \mathcal{O}_k^{A,+} \} \right) + c_{4,k}^A \text{Tr} \left(\overline{H}H \gamma_\mu \gamma_5 \mathbb{A}^\mu \right) \text{Tr}(\mathcal{O}_k^{A,+}) \\
& + c_{5,k}^A \text{Tr} \left(\overline{H}H \gamma_\mu \gamma_5 \right) \text{Tr}(\mathbb{A}^\mu \mathcal{O}_k^{A,+}) + c_{6,k}^A \text{Tr} \left(\overline{H}H \gamma_\mu [\mathbb{A}^\mu, \mathcal{O}_k^{A,-}] \right) \\
& + c_{7,k}^A \left(\text{Tr} \left(\overline{H}H \gamma_\mu \gamma_5 Q_k^A \mathbb{A}^\mu \tilde{Q}_k^A \right) + p.c. \right) \\
& \left. + c_{8,k}^A \left(\text{Tr} \left(\overline{H}H \gamma_\mu \gamma_5 Q_k^A \right) \text{Tr} \left(\mathbb{A}^\mu \tilde{Q}_k^A \right) + p.c. \right) \right\} \\
& + a^2 \sum_{k=2,5,7,8} c_{9,k}^A \left(\text{Tr} \left(\overline{H}H \gamma_\mu Q_k^A \mathbb{A}^\mu \tilde{Q}_k^A \right) + p.c. \right) \\
& + a^2 \sum_{k=1,2,6,7} c_{10,k}^A \left(\text{Tr} \left(\overline{H}H \gamma_\mu Q_k^A \right) \text{Tr} \left(\mathbb{A}^\mu \tilde{Q}_k^A \right) + p.c. \right). \tag{53}
\end{aligned}$$

Here, any taste indices in Q_k^A and $\tilde{Q}_k^A$ should be contracted; for example one should sum over λ and ν in a term involving $Q_6^A, \tilde{Q}_6^A$.⁵ Note that only a subset of the Q_k^A are allowed in the terms with coefficients $c_{9,k}^A$ and $c_{10,k}^A$, which involve γ_μ as opposed to $\gamma_\mu \gamma_5$. The parity conjugation introduces a minus sign, so the terms can only be Hermitian if $\tilde{Q}_k^{A\dagger} = Q_k^A$ ($c_{9,k}^A$ term) or $Q_k^{A\dagger} = Q_k^A$ ($c_{10,k}^A$ term). The forbidden terms are anti-Hermitian; adding a factor of i to make them Hermitian would violate $\mathcal{T}$ invariance.

The reader may wonder whether additional operators in $\mathcal{L}_{3,a^2}^A$ might be constructed using various Q_j^A and $\tilde{Q}_j^A$ in place of the $\mathcal{O}_k^{A,\pm}$. The answer is no, since the corresponding terms would contain either the product $Q_j^A \tilde{Q}_j^A$ or $Q_j^A \text{Tr}(\tilde{Q}_j^A)$ (along with appropriate parity-conjugated terms). In these circumstances, it is easy to check from Eq. (52) that one reproduces the $\mathcal{O}_k^{A,\pm}$.

The converse question is why the $Q_j^A, \tilde{Q}_j^A$ cannot be found by simply opening up a second Σ or $\Sigma^\dagger$ in the $\mathcal{O}_k^{A,\pm}$. The reason is that certain allowed combinations of spurions — namely $Q_2^A, \tilde{Q}_2^A$ and $Q_7^A, \tilde{Q}_7^A$ — have been eliminated from $\mathcal{V}_\Sigma$ and hence from $\mathcal{O}_k^{A,\pm}$ because their products are trivial and their traces vanish. Such combinations can only be found by returning to the original spurion analysis.

⁵ Contributions with $\lambda = \nu$ are automatically omitted since we use the definition $\xi_{\lambda\nu} = (1/2)[\xi_\lambda, \xi_\nu]$.

For the current at this order we cannot use $\mathbb{A}^\mu$, so the replacement $\mathcal{M}^\pm \rightarrow \mathcal{O}_k^{A,\pm}$ in Eq. (47) is all that is needed. We have

$$j_{2,a^2,A}^{\mu,b} = a^2 \sum_{k=1}^8 \left\{ r_{1,k}^A \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \mathcal{O}_k^{A,+} \sigma^\dagger \lambda^{(b)} + r_{2,k}^A \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} \text{Tr}(\mathcal{O}_k^{A,+}) \right. \\ \left. + r_{3,k}^A \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \mathcal{O}_k^{A,-} \sigma^\dagger \lambda^{(b)} + r_{4,k}^A \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} \text{Tr}(\mathcal{O}_k^{A,-}) \right\}. \quad (54)$$

B. Discretization errors at NLO: Operators from $\mathcal{L}_B^{\text{Sk}}$

The terms which arise from breaking rotation symmetry at this order can be determined through a spurion analysis that follows that of Sec. A3 in Ref. [30]. The following pairs of spurions can appear:

$$\begin{aligned} [V_\mu \times T_\mu], [A_\mu \times T_\mu] &\rightarrow Q_{\mu,1}^B = i\sigma \xi_{\mu\lambda}^{(n)} \sigma^\dagger, & \tilde{Q}_{\mu,1}^B &\equiv (Q_{\mu,1}^B)_{p.c.} = -i\sigma^\dagger \xi_{\lambda\mu}^{(n)} \sigma \\ [T_\mu \times V_\mu], &\rightarrow Q_{\mu,2}^B = \sigma \xi_\mu^{(n)} \sigma, & \tilde{Q}_{\mu,2}^B &\equiv (Q_{\mu,2}^B)_{p.c.} = \sigma^\dagger \xi_\mu^{(n)} \sigma^\dagger \\ [T_\mu \times A_\mu], &\rightarrow Q_{\mu,3}^B = i\sigma \xi_{\mu 5}^{(n)} \sigma, & \tilde{Q}_{\mu,3}^B &\equiv (Q_{\mu,3}^B)_{p.c.} = -i\sigma^\dagger \xi_{5\mu}^{(n)} \sigma^\dagger \\ [V_\mu \times T_\mu], [A_\mu \times T_\mu] &\rightarrow Q_{\mu,4}^B = i\sigma \xi_{\mu\lambda}^{(n)} \sigma^\dagger, & \tilde{Q}_{\mu,4}^B &\equiv Q_{\mu,4}^B, \end{aligned} \quad (55)$$

where we show the terms in $\mathcal{L}_B^{\text{Sk}}$ [Eq. (5)] that generate the given spurion operators. The index μ is singled out in the names $Q_{\mu,j}^B$ because it will be repeated 4 times in terms in $\mathcal{L}_{3,a^2}^B$, thereby breaking separate continuous rotations and taste symmetries, and leaving only the lattice symmetry of joint 90° rotations. We do not include the index λ in the names $Q_{\mu,1}^B$ and $Q_{\mu,4}^B$ because λ will appear only twice, and a sum over λ will be implied in any terms with $Q_{\mu,1}^B$, $\tilde{Q}_{\mu,1}^B$ or $Q_{\mu,4}^B$, $\tilde{Q}_{\mu,4}^B$. The spurions $Q_{\mu,j}^B$ ($j = 1, 2, 3, 4$) are in fact the same as Q_6^A , Q_5^A , Q_8^A and Q_7^A , respectively.

We can also construct operators, analogous to the $\mathcal{O}_k^{A,\pm}$, out of either $Q_{\mu,j}^B \tilde{Q}_{\mu,j}^B \pm p.c.$ or $Q_{\mu,j}^B \text{Tr}(\tilde{Q}_{\mu,j}^B) \pm p.c.$. These are

$$\begin{aligned} \mathcal{O}_{\mu,1}^{B,\pm} &= \sum_{\lambda \neq \mu} (\sigma \xi_{\mu\lambda}^{(n)} \Sigma^\dagger \xi_{\lambda\mu}^{(n)} \sigma) \pm p.c. , \\ \mathcal{O}_{\mu,2}^{B,\pm} &= (\sigma \xi_\mu^{(n)} \sigma) \text{Tr}(\xi_\mu^{(n)} \Sigma^\dagger) \pm p.c. , \\ \mathcal{O}_{\mu,3}^{B,\pm} &= (\sigma \xi_{\mu 5}^{(n)} \sigma) \text{Tr}(\xi_{5\mu}^{(n)} \Sigma^\dagger) \pm p.c. . \end{aligned} \quad (56)$$

The numbering here corresponds to that in Eq. (55); $Q_{\mu,4}^B$, $\tilde{Q}_{\mu,4}^B$ produce no operators of this kind. Note that the repeated index μ is not summed over in Eq. (56); aside from this these

operators are the same as operators $\mathcal{O}_6^{A,\pm}$, $\mathcal{O}_5^{A,\pm}$, and $\mathcal{O}_8^{A,\pm}$, respectively. Like the $\mathcal{O}_k^{A,\pm}$, the $\mathcal{O}_{\mu,k}^{B,\pm}$ transform according to Eq. (48).

To obtain terms that correspond to $\mathcal{L}_B^{\text{Sk}}$, we can multiply either the $\mathcal{O}_{\mu,k}^{B,\pm}$ or the $Q_{\mu,k}^B$, $\tilde{Q}_{\mu,k}^B$ by two additional four-vectors with index μ , and sum over μ . For $\mathcal{L}_{2,a^2}^B$, we can employ v_μ or γ_μ . We find:

$$\mathcal{L}_{2,a^2}^B = a^2 \sum_{\mu} \sum_{k=1}^3 \left\{ K_{1,k}^B v_\mu v^\mu \text{Tr}(\overline{H} H \mathcal{O}_{\mu,k}^{B,+}) + K_{2,k}^B v_\mu v^\mu \text{Tr}(\overline{H} H) \text{Tr}(\mathcal{O}_{\mu,k}^{B,+}) \right\} \quad (57)$$

We have used that fact $\text{Tr}(\overline{H} H \gamma_\mu) \propto \text{Tr}(\overline{H} H) v_\mu$ to eliminate other possibilities. Terms with $Q_{\mu,k}^B$, $\tilde{Q}_{\mu,k}^B$ are redundant and can be rewritten in terms of $\mathcal{O}_{\mu,k}^{B,\pm}$.

For $\mathcal{L}_{3,a^2}^B$, a single factor of D_μ or $\mathbb{A}_\mu$ is also allowed. The enumeration then gets quite complicated, because $Q_{\mu,k}^B$, $\tilde{Q}_{\mu,k}^B$ can appear in the $\mathbb{A}_\mu$ terms, and, in addition, various familiar simplifications cannot be used. For example, manipulations that follow from $H\psi = -H$ will not apply when the summation index μ in $\psi = v^\mu \gamma_\mu$ also appears in the taste-violating operators. To make the equations more manageable, we write,

$$\mathcal{L}_{3,a^2}^B = \mathcal{L}_{3,a^2}^{B,O} + \mathcal{L}_{3,a^2}^{B,Q} . \quad (58)$$

We then obtain:

$$\begin{aligned}
\mathcal{L}_{3,a^2}^{B,O} = a^2 \sum_{\mu} \sum_{k=1}^3 \Bigg\{ & i c_{1,k}^B \text{Tr} \left(\overline{H} H v^{\mu} \overleftarrow{D}_{\mu} \mathcal{O}_{\mu,k}^{B,+} - v^{\mu} \overrightarrow{D}_{\mu} \overline{H} H \mathcal{O}_{\mu,k}^{B,+} \right) \\
& + i c_{2,k}^B \text{Tr} \left(\overline{H} H v^{\mu} \overleftarrow{D}_{\mu} - v^{\mu} \overrightarrow{D}_{\mu} \overline{H} H \right) \text{Tr}(\mathcal{O}_{\mu,k}^{B,+}) \\
& + c_{3,k}^B \text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 \{ \mathbb{A}^{\mu}, \mathcal{O}_{\mu,k}^{B,+} \} \right) + c_{4,k}^B \text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 \mathbb{A}^{\mu} \right) \text{Tr}(\mathcal{O}_{\mu,k}^{B,+}) \\
& + c_{5,k}^B \text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 \right) \text{Tr}(\mathbb{A}^{\mu} \mathcal{O}_{\mu,k}^{B,+}) + c_{6,k}^B \text{Tr} \left(\overline{H} H \gamma_{\mu} [\mathbb{A}^{\mu}, \mathcal{O}_{\mu,k}^{B,-}] \right) \\
& + i c_{7,k}^B v_{\mu} v^{\mu} \text{Tr} \left(\overline{H} H v \cdot \overleftarrow{D} \mathcal{O}_{\mu,k}^{B,+} - v \cdot \overrightarrow{D} \overline{H} H \mathcal{O}_{\mu,k}^{B,+} \right) \\
& + i c_{8,k}^B v_{\mu} v^{\mu} \text{Tr} \left(\overline{H} H v \cdot \overleftarrow{D} - v \cdot \overrightarrow{D} \overline{H} H \right) \text{Tr}(\mathcal{O}_{\mu,k}^{B,+}) \\
& + c_{9,k}^B v_{\mu} v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\nu} \gamma_5 \{ \mathbb{A}^{\nu}, \mathcal{O}_{\mu,k}^{B,+} \} \right) + c_{10,k}^B v_{\mu} v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\nu} \gamma_5 \mathbb{A}^{\nu} \right) \text{Tr}(\mathcal{O}_{\mu,k}^{B,+}) \\
& + c_{11,k}^B v_{\mu} v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\nu} \gamma_5 \right) \text{Tr}(\mathbb{A}^{\nu} \mathcal{O}_{\mu,k}^{B,+}) + c_{12,k}^B v_{\mu} v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\nu} [\mathbb{A}^{\nu}, \mathcal{O}_{\mu,k}^{B,-}] \right) \\
& + c_{13,k}^B v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 \{ v \cdot \mathbb{A}, \mathcal{O}_{\mu,k}^{B,+} \} \right) + c_{14,k}^B v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 v \cdot \mathbb{A} \right) \text{Tr}(\mathcal{O}_{\mu,k}^{B,+}) \\
& + c_{15,k}^B v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 \right) \text{Tr}(v \cdot \mathbb{A} \mathcal{O}_{\mu,k}^{B,+}) + c_{16,k}^B v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\mu\nu} \gamma_5 \{ \mathbb{A}^{\nu}, \mathcal{O}_{\mu,k}^{B,+} \} \right) \\
& + c_{17,k}^B v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\mu\nu} \gamma_5 \right) \text{Tr} \left(\mathbb{A}^{\nu} \mathcal{O}_{\mu,k}^{B,+} \right) + c_{18,k}^B v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\mu\nu} \gamma_5 \mathbb{A}^{\nu} \right) \text{Tr} \left(\mathcal{O}_{\mu,k}^{B,+} \right) \\
& + c_{19,k}^B v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\mu\nu} \{ \mathbb{A}^{\nu}, \mathcal{O}_{\mu,k}^{B,-} \} \right) + c_{20,k}^B v^{\mu} \text{Tr} \left(\overline{H} H \gamma_{\mu\nu} \right) \text{Tr} \left(\mathbb{A}^{\nu} \mathcal{O}_{\mu,k}^{B,-} \right) \Bigg\} \quad (59)
\end{aligned}$$

A sum over the repeated index ν is implied, and $\gamma_{\mu\nu} \equiv (1/2)[\gamma_{\mu}, \gamma_{\nu}]$. We have used the fact that $\text{Tr}(\mathcal{O}_{\mu,k}^{B,-})$ vanishes for all k .

For $\mathcal{L}_{3,a^2}^{B,Q}$, we find:

$$\begin{aligned}
\mathcal{L}_{3,a^2}^{B,Q} = & a^2 \sum_{\mu} \left\{ \sum_{k=1}^4 \left[c_{21,k}^B \left(\text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 Q_{\mu,k}^B \mathbb{A}^{\mu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \right. \right. \\
& + c_{22,k}^B \left(\text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 Q_{\mu,k}^B \right) \text{Tr} \left(\mathbb{A}^{\mu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \\
& + c_{23,k}^B v_{\mu} v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\nu} \gamma_5 Q_{\mu,k}^B \mathbb{A}^{\nu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \\
& + c_{24,k}^B v_{\mu} v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\nu} \gamma_5 Q_{\mu,k}^B \right) \text{Tr} \left(\mathbb{A}^{\nu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \\
& + c_{25,k}^B v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 Q_{\mu,k}^B v \cdot \mathbb{A} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \\
& + c_{26,k}^B v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\mu} \gamma_5 Q_{\mu,k}^B \right) \text{Tr} \left(v \cdot \mathbb{A} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \\
& + c_{27,k}^B v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\mu\nu} \gamma_5 Q_{\mu,k}^B \mathbb{A}^{\nu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \\
& \left. + c_{28,k}^B v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\mu\nu} \gamma_5 Q_{\mu,k}^B \right) \text{Tr} \left(\mathbb{A}^{\nu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \right] \\
& + \sum_{k=2,3,4} \left[c_{29,k}^B \left(\text{Tr} \left(\overline{H} H \gamma_{\mu} Q_{\mu,k}^B \mathbb{A}^{\mu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \right. \\
& \left. + c_{30,k}^B v_{\mu} v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\nu} Q_{\mu,k}^B \mathbb{A}^{\nu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \right] \\
& + \sum_{k=1,4} \left[c_{31,k}^B \left(\text{Tr} \left(\overline{H} H \gamma_{\mu} Q_{\mu,k}^B \right) \text{Tr} \left(\mathbb{A}^{\mu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \right. \\
& \left. + c_{32,k}^B v_{\mu} v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\nu} Q_{\mu,k}^B \right) \text{Tr} \left(\mathbb{A}^{\nu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \right] \\
& + c_{33,1}^B v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\mu\nu} Q_{\mu,1}^B \mathbb{A}^{\nu} \tilde{Q}_{\mu,1}^B \right) + p.c. \right) \\
& \left. + \sum_{k=2,3} \left[c_{34,k}^B v^{\mu} \left(\text{Tr} \left(\overline{H} H \gamma_{\mu\nu} Q_{\mu,k}^B \right) \text{Tr} \left(\mathbb{A}^{\nu} \tilde{Q}_{\mu,k}^B \right) + p.c. \right) \right] \right\} \quad (60)
\end{aligned}$$

The additional implicit taste index λ that appears in $Q_{\mu,1}^B$, $\tilde{Q}_{\mu,1}^B$ or $Q_{\mu,4}^B$, $\tilde{Q}_{\mu,4}^B$ should be summed. As in Eq. (53), only a subset of the $Q_{\mu,k}^B$ are allowed by Hermiticity and $\mathcal{T}$ invariance in terms where the parity conjugate introduces a minus sign. Since $\mathbb{A}$ is an axial vector, these are the terms without an explicit γ_5 .

Note that we have raised one space-time μ index in each term in Eqs. (59) and (60) because we are using Minkowski space conventions. We also emphasize that the γ_{μ} here, and elsewhere in the chiral Lagrangian, are always Minkowski ones. The taste μ index on $\mathcal{O}_{\mu,k}^{B,\pm}$ and $Q_{\mu,k}^B$, $\tilde{Q}_{\mu,k}^B$ is not affected by going to Minkowski space, and taste matrices ξ_{μ} obey Euclidean conventions.

Proceeding to the current, we have:

$$\begin{aligned}
j_{2,a^2,B}^{\mu,b} = a^2 \sum_{k=1}^3 \bigg\{ & r_{1,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \mathcal{O}_{\mu,k}^{B,+} \sigma^\dagger \lambda^{(b)} + r_{2,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} \text{Tr}(\mathcal{O}_{\mu,k}^{B,+}) \\
& + r_{3,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \mathcal{O}_{\mu,k}^{B,-} \sigma^\dagger \lambda^{(b)} + r_{4,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} \text{Tr}(\mathcal{O}_{\mu,k}^{B,-}) \\
& + \sum_\nu \left(r_{5,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) v_\nu v^\nu \mathcal{O}_{\nu,k}^{B,+} \sigma^\dagger \lambda^{(b)} \right. \\
& \quad + r_{6,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} v_\nu v^\nu \text{Tr}(\mathcal{O}_{\nu,k}^{B,+}) \\
& \quad + r_{7,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) v_\nu v^\nu \mathcal{O}_{\nu,k}^{B,-} \sigma^\dagger \lambda^{(b)} \\
& \quad + r_{8,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H) \sigma^\dagger \lambda^{(b)} v_\nu v^\nu \text{Tr}(\mathcal{O}_{\nu,k}^{B,-}) \\
& \quad + r_{9,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H \gamma^\nu) v_\nu \mathcal{O}_{\nu,k}^{B,+} \sigma^\dagger \lambda^{(b)} \\
& \quad + r_{10,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H \gamma^\nu) \sigma^\dagger \lambda^{(b)} v_\nu \text{Tr}(\mathcal{O}_{\nu,k}^{B,+}) \\
& \quad + r_{11,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H \gamma^\nu) v_\nu \mathcal{O}_{\nu,k}^{B,-} \sigma^\dagger \lambda^{(b)} \\
& \quad \left. + r_{12,k}^B \text{tr}_D (\gamma^\mu (1 - \gamma_5) H \gamma^\nu) \sigma^\dagger \lambda^{(b)} v_\nu \text{Tr}(\mathcal{O}_{\nu,k}^{B,-}) \right) \bigg\}. \tag{61}
\end{aligned}$$

Here the repeated index μ is not summed over.

Despite the complexity of Eqs. (51) through (61), the effect of these terms on the heavy-light decay constant is extremely simple. At this order, the terms can contribute only at tree level, so we can set $\sigma = 1$ and $\mathbb{A}_\mu = 0$. Each term then either vanishes outright, is nonvanishing but does not contribute to f_B , or reduces to an additive constant in f_B that is proportional to a^2 . In particular, there are no rotation-violating contributions to f_B until higher order. All these a^2 analytic terms thus combine and lead merely to the presence of one additional unknown parameter in S χ PT chiral fits, compared to continuum-like chiral fits.

V. CHIRAL LOGARITHMS IN f_B AT ONE LOOP

Before calculating the one-loop diagrams, we need to write down the propagators of the heavy-light and pion fields, as well as the vertices coupling them. We use the notation $\{\psi\chi\}(k)$ to denote the Minkowski space propagator of fields ψ and χ with momentum k .

Then, from the LO heavy-light Lagrangian $\mathcal{L}_1$, Eq. (23), we have

$$\left\{ B_a B_b^\dagger \right\}(k) = \frac{i\delta_{ab}}{2(v \cdot k + i\epsilon)} , \quad (62)$$

$$\left\{ B_{\mu a}^* B_{\nu b}^{\dagger} \right\}(k) = \frac{-i\delta_{ab}(g_{\mu\nu} - v_\mu v_\nu)}{2(v \cdot k + i\epsilon)} . \quad (63)$$

Here a, b indicate the flavor-taste of the light quarks. Lower case Latin indices from the beginning of the alphabet ($a, b, c, d \dots$) will always serve as such flavor-taste indices; in the full QCD case they run over $4N_f$ values, where N_f is the number of light sea-quark flavors.

Since taste violations do not appear in $\mathcal{L}_1$, Eqs. (62) and (63) are the same as in the continuum theory [34], except that flavor-taste indices replace pure flavor indices. Similarly, the $BB^*\pi$ vertex looks identical to that in Ref. [34] aside from the redefinition of the indices and a factor of $1/2$ in the present case coming from our normalization of the taste generators, Eq. (15). The term in the interaction Lagrangian that gives this vertex is:

$$\frac{ig_\pi}{f} (B_{\mu a}^{\dagger} B_b - B_a^\dagger B_{\mu b}^*) \partial^\mu \Phi_{ba} , \quad (64)$$

where repeated indices are summed. The other needed vertex comes from the $B\pi\pi$ term in the expansion of the LO current, Eq. (24). We have:

$$j_{\text{LO}}^{\mu,b} = i\kappa v^\mu \left(B_b - \frac{1}{8f^2} B_a \Phi_{ac} \Phi_{cb} + \dots \right) , \quad (65)$$

where repeated indices are again summed and $\dots$ represents terms involving B^* or other numbers of pions, which do not contribute to the decay constant at this order.

When necessary, each flavor-taste index can be replaced by a pair of indices representing flavor and taste separately. We use Latin indices in the middle of the alphabet ($i, j, \dots$) as pure flavor indices, which take on the values $1, 2, \dots, N_f$ in full QCD. Greek indices at the beginning of the alphabet ($\alpha, \beta, \gamma, \dots$) will be used for quark taste indices, running from 1 to 4. Thus we can replace $a \rightarrow i\alpha$ and write, for example,

$$\left\{ B_{i\alpha} B_{j\beta}^\dagger \right\}(k) = \frac{i\delta_{ij}\delta_{\alpha\beta}}{2(v \cdot k + i\epsilon)} . \quad (66)$$

For light mesons, the LO Lagrangian $\mathcal{L}_{\text{LO}}^{\text{pions}}$, Eq. (27), includes both the continuum terms $\mathcal{L}_{\text{LO,cont}}^{\text{pions}}$ and the leading a^2 corrections $\mathcal{L}_{\text{LO},a^2}^{\text{pions}}$. As in Ref. [13], we treat explicitly the “hair-pin” terms arising from the m_0^2 contribution to Eq. (22) and the C_{2V} , C_{2A} , C_{5V} , and C_{5A}

contributions to Eq. (35) by separating out the disconnected parts of meson propagators. The remainder of $\mathcal{L}_{\text{LO}}^{\text{pions}}$ determines the connected pion propagators:

$$\left\{ \Phi_{ij}^{\Xi} \Phi_{j'i'}^{\Xi'} \right\}_{\text{conn}}(p) = \frac{i\delta_{ii'}\delta_{jj'}\delta_{\Xi\Xi'}}{p^2 - m_{ij,\Xi}^2 + i\epsilon} \quad (67)$$

where Ξ is one of the 16 meson tastes [as defined after Eq. (14)], and $m_{ij,\Xi}$ is the tree-level mass of a taste- Ξ meson composed of quarks of flavor i and j :

$$m_{ij,\Xi}^2 = \mu(m_i + m_j) + a^2\Delta_{\Xi}. \quad (68)$$

Here Δ_{Ξ} is the taste splitting, which can be expressed in terms of C_1 , C_3 , C_4 and C_6 in Eq. (35) [13].

Because of the residual $SO(4)$ taste symmetry [11] at this order, the mesons within a given taste multiplet (P , V , T , A , or I) are degenerate in mass. When it is not important to specify the particular member of the multiplet in question, we will usually just name the multiplet, for example:

$$m_{ij,V}^2 = \mu(m_i + m_j) + a^2\Delta_V. \quad (69)$$

Since the heavy-light propagators are most simply written with flavor-taste indices, as in Eqs. (62) and (63), it is convenient for current purposes to rewrite Eq. (67) in flavor-taste notation also:

$$\left\{ \Phi_{ab} \Phi_{b'a'} \right\}_{\text{conn}}(p) \equiv \left\{ \Phi_{i\alpha,j\beta} \Phi_{j'\beta',i'\alpha'} \right\}_{\text{conn}}(p) = \sum_{\Xi} \frac{i\delta_{ii'}\delta_{jj'}T_{\alpha\beta}^{\Xi}T_{\beta'\alpha'}^{\Xi}}{p^2 - m_{ij,\Xi}^2 + i\epsilon}, \quad (70)$$

where T^{Ξ} are the 16 taste generators, Eq. (15).

For flavor-charged pions ($i \neq j$), the complete propagators are just the connected propagators in Eq. (67) or (70). However, for flavor-neutral pions ($i = j$), there are disconnected contributions coming from one or more hairpin insertions. At LO, these appear only for taste singlet, vector, or axial-vector pions. Denoting the Minkowski hairpin vertices as $-i\delta'_{\Xi}$, we have [13]:

$$\delta'_{\Xi} = \begin{cases} a^2\delta'_V, & T_{\Xi} \in \{\xi_{\mu}\} \text{ (taste vector);} \\ a^2\delta'_A, & T_{\Xi} \in \{\xi_{\mu 5}\} \text{ (taste axial-vector);} \\ 4m_0^2/3, & T_{\Xi} = \xi_I \text{ (taste singlet);} \\ 0, & T_{\Xi} \in \{\xi_{\mu\nu}, \xi_5\} \text{ (taste tensor or pseudoscalar)} \end{cases} \quad (71)$$

with

$$\delta'_{V(A)} \equiv \frac{16}{f^2}(C_{2V(A)} - C_{5V(A)}) . \quad (72)$$

The disconnected pion propagator is then

$$\left\{ \Phi_{ij}^{\Xi} \Phi_{j'i'}^{\Xi'} \right\}_{\text{disc}}(p) = \delta_{ij} \delta_{j'i'} \delta_{\Xi\Xi'} \mathcal{D}_{ii,i'i'}^{\Xi} , \quad (73)$$

where [13]

$$\mathcal{D}_{ii,i'i'}^{\Xi} = -i\delta'_{\Xi} \frac{i}{(p^2 - m_{ii,\Xi}^2)} \frac{i}{(p^2 - m_{i'i',\Xi}^2)} \frac{(p^2 - m_{U,\Xi}^2)(p^2 - m_{D,\Xi}^2)(p^2 - m_{S,\Xi}^2)}{(p^2 - m_{\pi^0,\Xi}^2)(p^2 - m_{\eta,\Xi}^2)(p^2 - m_{\eta',\Xi}^2)} . \quad (74)$$

For concreteness we have assumed that there are three sea-quark flavors: u , d , and s ; the generalization to N_f flavors is immediate. Here $m_{U,\Xi} \equiv m_{uu,\Xi}$ is the mass of a taste- Ξ pion made from a u and a $\bar{u}$ quark, neglecting hairpin mixing (and similarly for $m_{D,\Xi}$ and $m_{S,\Xi}$), $m_{\pi^0,\Xi}$, $m_{\eta,\Xi}$, and $m_{\eta',\Xi}$ are the mass eigenvalues after mixing is included, and the $i\epsilon$ terms have been left implicit. When specifying the particular member of a taste multiplet appearing in the disconnected propagator is unnecessary, we will abuse this notation slightly following Eq. (69) and refer to $\mathcal{D}_{ii,i'i'}^V$, $\mathcal{D}_{ii,i'i'}^A$, or $\mathcal{D}_{ii,i'i'}^I$. In flavor-taste notation we have:

$$\left\{ \Phi_{ab} \Phi_{b'a'} \right\}_{\text{disc}}(p) \equiv \left\{ \Phi_{i\alpha,j\beta} \Phi_{j'\beta',i'\alpha'} \right\}_{\text{disc}}(p) = \delta_{ij} \delta_{j'i'} \sum_{\Xi} T_{\alpha\beta}^{\Xi} T_{\beta'\alpha'}^{\Xi} \mathcal{D}_{ii,i'i'}^{\Xi} \quad (75)$$

Equations (73), (74), and (75) apply both to full and partially quenched QCD. In the latter case, i and i' can represent (quenched) valence quarks. Below, we will need the disconnected propagator only when i and i' both denote a particular valence quark: call it flavor x . Letting X be the flavor-neutral meson consisting of an x and an $\bar{x}$, we will use the simplified notation $\mathcal{D}_{XX}^{\Xi} \equiv \mathcal{D}_{xx,xx}^{\Xi}$ or, for example, $\mathcal{D}_{XX}^V \equiv \mathcal{D}_{xx,xx}^V$.

We now calculate the one-loop decay constant, f_{B_a} , of a heavy-light pseudoscalar meson with light flavor-taste a . The calculation is done in the 4+4+4 theory, where there are four tastes per flavor. We then adjust the result for the 1+1+1 (one taste per flavor) case of interest. We work out explicitly the partially quenched case; full theory results are easily obtained by taking appropriate limits. The valence flavor is called x ; the valence taste, α . In other words, we replace $a \rightarrow x\alpha$. As will be seen, the decay constant does not depend on the taste of the valence quark, so we may write $f_{B_a} \rightarrow f_{B_x}$.

At one loop, we express the decay constant as

$$f_{B_x} = f_{B_x}^{\text{LO}} \left(1 + \frac{1}{16\pi^2 f^2} \delta f_{B_x} + \text{analytic terms} \right) . \quad (76)$$

Recall that the lowest order term $f_{B_x}^{\text{LO}}$ depends on the light valence flavor in a rather trivial way: $f_{B_x}^{\text{LO}} = \kappa/\sqrt{m_{B_x}}$. The one-loop diagrams that contribute to δf_{B_x} are shown in Figs. 1 and 2, where the cross indicates a disconnected pion propagator (*i.e.*, one or more hairpin insertions for the singlet, vector and axial tastes). The terms in Fig. 3 vanish because the integrals produce an overall factor of v_ν on shell, which is then multiplied by the projector $g^{\mu\nu} - v^\mu v^\nu$.

The diagrams in Fig. 1 contribute to the heavy-light wavefunction renormalization. Using the propagators and vertices defined above, one easily sees that the self energy S_{ab} is proportional to the identity in taste space, and is diagonal in flavor. We thus have

$$S_{ab}(v \cdot k) \equiv S_{i\alpha,j\beta}(v \cdot k) = \delta_{\alpha\beta} \delta_{ij} \tilde{S}_i(v \cdot k) . \quad (77)$$

Flavor symmetry (broken by the diagonal mass terms) guarantees this flavor structure. The taste-independence follows from the residual discrete taste symmetry of the light quarks, which corresponds to the shift symmetry in the staggered action. This symmetry is

$$q_i \rightarrow (1 \otimes \xi_\mu) q_i , \quad \bar{q}_i \rightarrow \bar{q}_i (1 \otimes \xi_\mu) , \quad (78)$$

at the level of the Symanzik action; while at the chiral level it is

$$\begin{aligned} \Sigma &\rightarrow \xi_\mu^{(n)} \Sigma \xi_\mu^{(n)} , \\ \sigma &\rightarrow \xi_\mu^{(n)} \sigma \xi_\mu^{(n)} , \\ H &\rightarrow H \xi_\mu^{(n)} , \\ \overline{H} &\rightarrow \xi_\mu^{(n)} \overline{H} . \end{aligned} \quad (79)$$

Note that the symmetry is diagonal in flavor; the transformation acts only on the taste indices and affects all light quark flavors identically.

The one-loop wavefunction renormalization, δZ_{B_x} , for B_x is then

$$\delta Z_{B_x} = \frac{1}{2} \frac{d\tilde{S}_x(v \cdot k)}{d(v \cdot k)} \bigg|_{v \cdot k=0} . \quad (80)$$

This contributes an amount $\delta f_{B_x}^Z$ to δf_{B_x} . We find (in the 4+4+4 theory):

$$\begin{aligned} \delta f_{B_x}^Z &= \frac{1}{2} (16\pi^2 f^2) \delta Z_{B_x} \\ &= \frac{g_\pi^2}{8} (g_{\mu\nu} - v_\mu v_\nu) \int \frac{d^4 p}{\pi^2} \frac{p^\mu p^\nu}{(v \cdot p + i\epsilon)^2} \left[\sum_{f,\Xi} \frac{i}{p^2 - m_{xf,\Xi}^2 + i\epsilon} \right. \\ &\quad \left. + \mathcal{D}_{XX}^I + 4\mathcal{D}_{XX}^V + 4\mathcal{D}_{XX}^A \right] \end{aligned} \quad (81)$$

where Ξ runs over the 16 pion tastes, and f runs over the three sea quarks, u , d , and s . The term summed over f and Ξ arises from the “connected” diagram, Fig. 1(a); as we will see below, it involves an internal sea quark loop. The remaining terms come from the “disconnected” diagram, Fig. 1(b). We have summed over the four degenerate vector and axial disconnected contributions, as seen by the factors of four in front of $\mathcal{D}^V$ and $\mathcal{D}^A$.

The current corrections contribute an amount $\delta f_{B_x}^{\text{cur}}$ to δf_{B_x} . Computing the diagrams in Fig. 2, we obtain (in the 4+4+4 theory):

$$\delta f_{B_x}^{\text{cur}} = -\frac{1}{8} \int \frac{d^4 p}{\pi^2} \left[\sum_{f,\Xi} \frac{i}{p^2 - m_{xf,\Xi}^2 + i\epsilon} + \mathcal{D}_{XX}^I + 4\mathcal{D}_{XX}^V + 4\mathcal{D}_{XX}^A \right]. \quad (82)$$

These chiral integrals are similar to those that appear [13] for the decay constant, f_P , of a partially quenched pion P . The main difference is that here we have a single light valence quark x , whereas P is a bound state of two light valence quarks, x and $\bar{y}$. So the corresponding integrals in the f_P case have additional contributions with $x \rightarrow y$ in various terms.

We now need to convert Eqs. (81) and (82) from a 4+4+4 to a 1+1+1 theory. This can easily be done with a quark-flow analysis, following Ref. [13]. The g_π vertex at the quark level is shown in Fig. 4(a); the current vertex at second order in the pion fields is depicted in Fig. 4(b). The connected terms in Eqs. (81) and (82) that come from these vertices thus involve an internal quark loop, shown in Fig. 5(a), and should be multiplied by a factor of 1/4. As described in Ref. [13], the only other changes in going from 4+4+4 to 1+1+1 appear in the mass eigenstates of the full flavor-neutral propagators. This is due to the factors of 1/4 that are associated with iteration of the two-point vertex, Eq. (71), in Fig. 5(c).

In the 1+1+1 theory, we thus have

$$\delta f_{B_x}^Z = \frac{g_\pi^2}{8} (g_{\mu\nu} - v_\mu v_\nu) \int \frac{d^4 p}{\pi^2} \frac{p^\mu p^\nu}{(v \cdot p + i\epsilon)^2} \left[\frac{1}{4} \sum_{f,\Xi} \frac{i}{p^2 - m_{xf,\Xi}^2 + i\epsilon} + \mathcal{D}_{XX}^I + 4\mathcal{D}_{XX}^V + 4\mathcal{D}_{XX}^A \right] \quad (83)$$

$$\delta f_{B_x}^{\text{cur}} = -\frac{1}{8} \int \frac{d^4 p}{\pi^2} \left[\frac{1}{4} \sum_{f,\Xi} \frac{i}{p^2 - m_{xf,\Xi}^2 + i\epsilon} + \mathcal{D}_{XX}^I + 4\mathcal{D}_{XX}^V + 4\mathcal{D}_{XX}^A \right], \quad (84)$$

where the mass eigenvalues that appear in $\mathcal{D}^I$, $\mathcal{D}^V$, and $\mathcal{D}^A$ [c.f. Eq. (74)] are now the 1+1+1 values [13]. For example, with three sea-quark flavors, $m_{\eta',I} \sim m_0^2$ for large m_0^2 , rather than

the value $4m_0^2$ that follows from Eq. (22). After the $m_0^2 \rightarrow \infty$ limit is taken, this implies that the $\mathcal{D}^I$ contribution is 4 times larger in the 1+1+1 theory than in the 4+4+4 theory.

The disconnected propagators in Eqs. (83) and (84) can then be written as a sum of single or double poles using the residue functions introduced in Ref. [13]. We define $\{m\} \equiv \{m_1, m_2, \dots, m_n\}$ as the set of masses that appear in the denominator of Eq. (74), and $\{\mu\} \equiv \{\mu_1, \mu_2, \dots, \mu_k\}$ as the numerator set of masses. Then, for $n > k$ and all masses distinct, we have:

$$\mathcal{I}^{[n,k]}(\{m\};\{\mu\}) \equiv \frac{\prod_{i=1}^k (q^2 - \mu_i^2)}{\prod_{j=1}^n (q^2 - m_j^2 + i\epsilon)} = \sum_{j=1}^n \frac{(-1)^{n+k+1} R_j^{[n,k]}(\{m\};\{\mu\})}{q^2 - m_j^2 + i\epsilon}, \quad (85)$$

where

$$R_j^{[n,k]}(\{m\};\{\mu\}) \equiv \frac{\prod_{i=1}^k (\mu_i^2 - m_j^2)}{\prod_{r \neq j} (m_r^2 - m_j^2)}. \quad (86)$$

If there is one double pole term for $q^2 = m_\ell^2$ (where $m_\ell \in \{m\}$), then

$$\begin{aligned} \mathcal{I}_{\text{dp}}^{[n,k]}(m_\ell; \{m\}; \{\mu\}) &\equiv \frac{\prod_{i=1}^k (q^2 - \mu_i^2)}{(q^2 - m_\ell^2 + i\epsilon) \prod_{j=1}^n (q^2 - m_j^2 + i\epsilon)} \\ &= \frac{\partial}{\partial m_\ell^2} \sum_{j=1}^n \frac{(-1)^{n+k+1} R_j^{[n,k]}(\{m\};\{\mu\})}{q^2 - m_j^2 + i\epsilon}. \end{aligned} \quad (87)$$

We use the same definitions of the residues $R_j^{[n,k]}$ as in Ref. [13]; the factors of $(-1)^{n+k+1}$ arise from the fact that the residues were defined for Euclidean space integrals, but we are working in Minkowski space here.

Using Eqs. (85) and (87), the integration in Eq. (84) is immediate, and that in Eq. (83) is standard in the continuum heavy-light literature. We follow Ref. [36] and define (in $4 - \hat{\epsilon}$ dimensions, with chiral scale Λ_χ)

$$\begin{aligned} J^{\mu\nu}(m, \Delta) &\equiv i\Lambda_\chi^{\hat{\epsilon}} \int \frac{d^{4-\hat{\epsilon}}p}{(2\pi)^{4-\hat{\epsilon}}} \frac{p^\mu p^\nu}{(p^2 - m^2 + i\epsilon)(v \cdot p - \Delta + i\epsilon)} \\ &= \frac{1}{16\pi^2} \Delta [J_1(m, \Delta) g^{\mu\nu} + J_2(m, \Delta) v^\mu v^\nu]. \end{aligned} \quad (88)$$

Δ in this equation should not be confused with the Δ_Ξ defined in Eq. (68). For Eq. (83), we need only

$$(g_{\mu\nu} - v_\mu v_\nu) \frac{\partial J^{\mu\nu}(m, \Delta)}{\partial \Delta} \Big|_{\Delta=0} = \frac{3}{16\pi^2} \frac{\partial \Delta J_1(m, \Delta)}{\partial \Delta} \Big|_{\Delta=0} \rightarrow -\frac{3}{16\pi^2} \ell(m^2) \quad (89)$$

$$(g_{\mu\nu} - v_\mu v_\nu) \frac{\partial}{\partial m^2} \frac{\partial J^{\mu\nu}(m, \Delta)}{\partial \Delta} \Big|_{\Delta=0} = \frac{3}{16\pi^2} \frac{\partial}{\partial m^2} \frac{\partial J_1(m, \Delta)}{\partial \Delta} \Big|_{\Delta=0} \rightarrow \frac{3}{16\pi^2} \tilde{\ell}(m^2) \quad (90)$$

As in Ref. [13], we use the arrow to denote the fact that we drop the analytic terms and keep only the chiral logarithms. The functions ℓ and $\tilde{\ell}$ are defined as

$$\ell(m^2) \equiv m^2 \ln \frac{m^2}{\Lambda_\chi^2} \quad [\text{infinite volume}] , \quad (91)$$

$$\tilde{\ell}(m^2) \equiv -\frac{\partial}{\partial m^2} \ell(m^2) = -\ln \left(\frac{m^2}{\Lambda_\chi^2} \right) - 1 \quad [\text{infinite volume}] . \quad (92)$$

These are the infinite volume forms; the effect of finite spatial volume will be discussed in Sec. VIII.

For Eq. (84) we need the Minkowski version of the integrals defined in Ref. [13]:

$$I_1(m^2) \equiv i\Lambda_\chi^\epsilon \int \frac{d^{4-\epsilon}p}{(2\pi)^{4-\epsilon}} \frac{1}{(p^2 - m^2 + i\epsilon)} \rightarrow \frac{1}{16\pi^2} \ell(m^2) \quad (93)$$

$$I_2(m^2) \equiv i\Lambda_\chi^\epsilon \int \frac{d^{4-\epsilon}p}{(2\pi)^{4-\epsilon}} \frac{1}{(p^2 - m^2 + i\epsilon)^2} \rightarrow -\frac{1}{16\pi^2} \tilde{\ell}(m^2) . \quad (94)$$

VI. FINAL NLO RESULTS

Now we are ready to write down the full NLO result, including the analytic terms.

After performing the integrals using Eqs. (89) through (94), we get for the 1+1+1 partially quenched case with all masses unequal:

$$\begin{aligned} \left(\frac{f_{B_x}}{f_{B_x}^{\text{LO}}} \right)_{1+1+1} = & 1 + \frac{1}{16\pi^2 f^2} \frac{1+3g_\pi^2}{2} \left\{ -\frac{1}{16} \sum_{f,\Xi} \ell(m_{xf,\Xi}^2) \right. \\ & - \frac{1}{3} \sum_{j \in \mathcal{M}_I^{(3,x)}} \frac{\partial}{\partial m_{X,I}^2} \left[R_j^{[3,3]}(\mathcal{M}_I^{(3,x)}; \mu_I^{(3)}) \ell(m_j^2) \right] \\ & - \left(a^2 \delta'_V \sum_{j \in \mathcal{M}_V^{(4,x)}} \frac{\partial}{\partial m_{X,V}^2} \left[R_j^{[4,3]}(\mathcal{M}_V^{(4,x)}; \mu_V^{(3)}) \ell(m_j^2) \right] + [V \rightarrow A] \right) \Big\} \\ & + c_s(m_u + m_d + m_s) + c_v m_x + c_a a^2 \end{aligned} \quad (95)$$

where f runs over the three sea quarks u , d , and s , Ξ runs over the 16 meson tastes, and the sets of masses in the residues are defined as follows (with taste labels implicit):

$$\mu^{(3)} = \{m_U^2, m_D^2, m_S^2\} , \quad (96)$$

$$\mathcal{M}^{(3,x)} = \{m_X^2, m_{\pi^0}^2, m_\eta^2\} , \quad (97)$$

$$\mathcal{M}^{(4,x)} = \{m_X^2, m_{\pi^0}^2, m_\eta^2, m_{\eta'}^2\} . \quad (98)$$

The analytic terms in Eq. (95) come from wave function renormalization contributions from $\mathcal{L}_3$ [Eqs. (46), (53) and (58)], as well as from higher order corrections to the current $j_2^{\mu,b}$ [Eqs. (47), (54) and (61)]. The low energy constants (LECs) c_s , c_v and c_a can be written as linear combinations of the large number of constants appearing in $\mathcal{L}_3$ and $j_2^{\mu,b}$. However, such expressions are unlikely to be useful, and we omit them here.

Of relevance to MILC simulations is the case where the up and down quark masses are degenerate, the 2+1 case. Setting $m_u = m_d = m_l$, we obtain

$$\begin{aligned} \left(\frac{f_{B_x}}{f_{B_x}^{\text{LO}}} \right)_{2+1} = & 1 + \frac{1}{16\pi^2 f^2} \frac{1+3g_\pi^2}{2} \left\{ -\frac{1}{16} \sum_{f,\Xi} \ell(m_{xf,\Xi}^2) \right. \\ & - \frac{1}{3} \sum_{j \in \mathcal{M}_I^{(2,x)}} \frac{\partial}{\partial m_{X,I}^2} \left[R_j^{[2,2]}(\mathcal{M}_I^{(2,x)}; \mu_I^{(2)}) \ell(m_j^2) \right] \\ & - \left(a^2 \delta'_V \sum_{j \in \hat{\mathcal{M}}_V^{(3,x)}} \frac{\partial}{\partial m_{X,V}^2} \left[R_j^{[3,2]}(\hat{\mathcal{M}}_V^{(3,x)}; \mu_V^{(2)}) \ell(m_j^2) \right] + [V \rightarrow A] \right) \Big\} \\ & + c_s(2m_l + m_s) + c_v m_x + c_a a^2, \end{aligned} \quad (99)$$

where

$$\mu^{(2)} = \{m_U^2, m_S^2\}, \quad (100)$$

$$\mathcal{M}^{(2,x)} = \{m_X^2, m_\eta^2\}, \quad (101)$$

$$\hat{\mathcal{M}}^{(3,x)} = \{m_X^2, m_\eta^2, m_{\eta'}^2\}. \quad (102)$$

Another interesting limit is the case of N_f degenerate flavors of sea quarks of mass m_f .

$$\begin{aligned} \left(\frac{f_{B_x}}{f_{B_x}^{\text{LO}}} \right)_{N_f \text{ degen}} = & 1 + \frac{1}{16\pi^2 f^2} \frac{1+3g_\pi^2}{2} \left\{ -\frac{N_f}{16} \sum_{\Xi} \ell(m_{xf,\Xi}^2) \right. \\ & - \frac{1}{N_f} \left[(m_{X,I}^2 - m_{ff,I}^2) \tilde{\ell}(m_{X,I}^2) - \ell(m_{X,I}^2) \right] \\ & - \left(a^2 \delta'_V \frac{\partial}{\partial m_{X,V}^2} \left[\frac{(m_{ff,V}^2 - m_{X,V}^2) \ell(m_{X,V}^2) - (m_{ff,V}^2 - m_{\eta',V}^2) \ell(m_{\eta',V}^2)}{m_{\eta',V}^2 - m_{X,V}^2} \right] \right. \\ & \left. \left. + [V \rightarrow A] \right) \right\} + N_f c_s m_f + c_v m_x + c_a a^2, \end{aligned} \quad (103)$$

where, in this case,

$$m_{\eta',V}^2 = m_{ff,V}^2 + \frac{N_f}{4} a^2 \delta'_V, \quad (104)$$

and similarly for $m_{\eta',A}^2$. The result in Eq. (103) reduces to that of Sharpe and Zhang [38] in the continuum limit.⁶ The $1/N_f$ dependence of the taste-singlet term in Eq. (103) is characteristic of the disconnected contribution and arises from the fact that $m_{\eta',I} \sim N_f m_0^2/3$ for large m_0^2 in the $1+1+1+\dots$ theory.

We now turn to full QCD, where the valence quark is one of the light sea quarks. In the $2+1$ case, we have

$$\begin{aligned} \left(\frac{f_{B_u}}{f_{B_u}^{\text{LO}}} \right)_{2+1 \text{ full}} &= 1 + \frac{1}{16\pi^2 f^2} \left(\frac{1+3g_\pi^2}{2} \right) \left\{ -\frac{1}{16} \sum_{\Xi} [2\ell(m_{\pi,\Xi}^2) + \ell(m_{K,\Xi}^2)] \right. \\ &\quad + \frac{1}{2} \ell(m_{\pi,I}^2) - \frac{1}{6} \ell(m_{\eta,I}^2) \\ &\quad - \left(a^2 \delta'_V \left[\frac{(m_{\pi,V}^2 - m_{S,V}^2)}{(m_{\pi,V}^2 - m_{\eta,V}^2)(m_{\pi,V}^2 - m_{\eta',V}^2)} \ell(m_{\pi,V}^2) \right. \right. \\ &\quad + \frac{(m_{\eta,V}^2 - m_{S,V}^2)}{(m_{\eta,V}^2 - m_{\pi,V}^2)(m_{\eta,V}^2 - m_{\eta',V}^2)} \ell(m_{\eta,V}^2) \\ &\quad \left. \left. + \frac{(m_{\eta',V}^2 - m_{S,V}^2)}{(m_{\eta',V}^2 - m_{\pi,V}^2)(m_{\eta',V}^2 - m_{\eta,V}^2)} \ell(m_{\eta',V}^2) \right] + [V \rightarrow A] \right) \Big\} \\ &\quad + c_s(2m_l + m_s) + c_v m_l + c_a a^2, \end{aligned} \quad (105)$$

$$\begin{aligned} \left(\frac{f_{B_s}}{f_{B_s}^{\text{LO}}} \right)_{2+1 \text{ full}} &= 1 + \frac{1}{16\pi^2 f^2} \left(\frac{1+3g_\pi^2}{2} \right) \left\{ -\frac{1}{16} \sum_{\Xi} [\ell(m_{S,\Xi}^2) + 2\ell(m_{K,\Xi}^2)] \right. \\ &\quad + \ell(m_{S,I}^2) - \frac{2}{3} \ell(m_{\eta,I}^2) \\ &\quad - \left(a^2 \delta'_V \left[\frac{(m_{S,V}^2 - m_{\pi,V}^2)}{(m_{S,V}^2 - m_{\eta,V}^2)(m_{S,V}^2 - m_{\eta',V}^2)} \ell(m_{S,V}^2) \right. \right. \\ &\quad + \frac{(m_{\eta,V}^2 - m_{\pi,V}^2)}{(m_{\eta,V}^2 - m_{S,V}^2)(m_{\eta,V}^2 - m_{\eta',V}^2)} \ell(m_{\eta,V}^2) \\ &\quad \left. \left. + \frac{(m_{\eta',V}^2 - m_{\pi,V}^2)}{(m_{\eta',V}^2 - m_{S,V}^2)(m_{\eta',V}^2 - m_{\eta,V}^2)} \ell(m_{\eta',V}^2) \right] + [V \rightarrow A] \right) \Big\} \\ &\quad + c_s(2m_l + m_s) + c_v m_s + c_a a^2. \end{aligned} \quad (106)$$

We have used the fact that $m_{\eta,I}^2 = (2m_{S,I}^2 + m_{\pi,I}^2)/3$ at this order to simplify the taste-singlet residues. Note that, as expected for full QCD, there are no longer any double-pole contributions. In the continuum limit Eqs. (105) and (106) reduce to known full QCD results [32, 33, 38].

⁶ Note that the analytic term coming from the -1 in our Eq. (92) is absorbed into the LECs in Eq. (3.6) of Ref. [38].

VII. CHIRAL-SCALE DEPENDENCE AND LOW ENERGY CONSTANTS

The effect of a change in the chiral scale Λ_χ can be absorbed into the LECs c_s , c_v , and c_a . In fact, the change of most of the one-loop terms is proportional to taste-split squared meson masses, rather than quark masses *per se*, since it is the meson masses that appear in the non-analytic terms. It has therefore proved convenient for light [10] and heavy-light [9] lattice calculations to rewrite the analytic terms as functions of meson masses, rather than quark masses, and, specifically, to use those combinations of meson masses that arise naturally from a change in scale in the non-analytic terms. Since squared meson masses have additive contributions proportional to a^2 due to taste violations [see Eq. (68)], this procedure automatically introduces some a^2 terms proportional to c_v and c_s , and is equivalent to a redefinition $c_a \rightarrow C_a = c_a - \beta_v c_v - \beta_s c_s$, where $\beta_{v,s}$ are constants.

The redefinition is convenient because the scale dependence of C_a is simplified and no longer depends on the splittings Δ_Ξ . More importantly, the procedure has a practical advantage for reducing lattice errors. From the simulations, C_a seems to be significantly smaller than c_a . In other words, most of the discretization error from the light quarks appears to be due to the a^2 dependence of the light meson masses in the chiral loops. To the extent this is true, *i.e.*, that C_a can be neglected in first approximation, the a^2 dependence is tied to the mass dependence through c_v and c_s . This means that a determination of the mass dependence at fixed lattice spacing can be used to estimate the lattice-spacing dependence.

The appropriate redefinitions do not depend on the quark masses, but, like the LECs themselves, they do depend on N_f . Therefore it is most convenient to work with the case of degenerate sea quark masses but arbitrary N_f , Eq. (103). Using Eq. (68), we can write the change in the chiral log contribution δf_{B_x} [*c.f.* Eq. (76)] under a change $\Lambda_\chi \rightarrow \tilde{\Lambda}_\chi$ in the chiral scale as

$$\begin{aligned} \delta f_{B_x} \rightarrow \delta f_{B_x} + \log(\tilde{\Lambda}^2/\Lambda^2) \frac{(1 + 3g_\pi^2)}{2} \left\{ \frac{N_f^2 - 4}{2N_f} \left(2\mu m_x + a^2 \Delta_{\text{val}} \right) \right. \\ \left. + \frac{N_f^2 + 2}{2N_f} \left(2\mu m_f + a^2 \Delta_{\text{sea}} \right) + a^2 (\delta'_V + \delta'_A) \right\}, \quad (107) \end{aligned}$$

where

$$\Delta_{\text{val}} \equiv \frac{N_f^2}{N_f^2 - 4} \bar{\Delta} - \frac{4}{N_f^2 - 4} \Delta_I \quad (108)$$

$$\Delta_{\text{sea}} \equiv \frac{N_f^2}{N_f^2 + 2} \bar{\Delta} + \frac{2}{N_f^2 + 2} \Delta_I , \quad (109)$$

with Δ_I the taste-singlet splitting and $\bar{\Delta}$ the average splitting:

$$\bar{\Delta} = \frac{1}{16} \sum_{\Xi} \Delta_{\Xi} = \frac{1}{16} (4\Delta_A + 6\Delta_T + 4\Delta_V + \Delta_I) . \quad (110)$$

Based on Eq. (107), it is natural to redefine the LECs by

$$N_f c_s m_f + c_v m_x + c_a a^2 \rightarrow N_f C_s (2\mu m_f + a^2 \Delta_{\text{sea}}) + C_v (2\mu m_x + a^2 \Delta_{\text{val}}) + C_a a^2 \quad (111)$$

for degenerate quarks [*e.g.*, in Eq. (103)], or

$$\begin{aligned} c_s(m_u + m_d + m_s + \dots) + c_v m_x + c_a a^2 &\rightarrow C_s (2\mu (m_u + m_d + m_s + \dots) + N_f a^2 \Delta_{\text{sea}}) \\ &\quad + C_v (2\mu m_x + a^2 \Delta_{\text{val}}) + C_a a^2 \end{aligned} \quad (112)$$

in the general case [*e.g.*, in Eq. (95)]. With these redefinitions, the change in the LECs under a chiral scale change is rather simple:

$$\begin{aligned} C_s(\tilde{\Lambda}_\chi) &= C_s(\Lambda_\chi) - \log(\tilde{\Lambda}_\chi^2/\Lambda_\chi^2) \frac{(1 + 3g_\pi^2)}{32\pi^2 f^2} \frac{(N_f^2 + 2)}{2N_f^2} \\ C_v(\tilde{\Lambda}_\chi) &= C_v(\Lambda_\chi) - \log(\tilde{\Lambda}_\chi^2/\Lambda_\chi^2) \frac{(1 + 3g_\pi^2)}{32\pi^2 f^2} \frac{(N_f^2 - 4)}{2N_f} \\ C_a(\tilde{\Lambda}_\chi) &= C_a(\Lambda_\chi) - \log(\tilde{\Lambda}_\chi^2/\Lambda_\chi^2) \frac{(1 + 3g_\pi^2)}{32\pi^2 f^2} (\delta'_v + \delta'_A) . \end{aligned} \quad (113)$$

We emphasize that the redistribution of the a^2 terms in Eq. (111) or (112) is not unique, and it is possible to move them from C_s to C_v without complicating the transformation properties of C_a under chiral scale change. In the “mixed” meson terms $m_{xf,\Xi}$ in Eq. (103), for example, we have chosen to associate half of each splitting with the valence quark x and half with the sea quark f . This is simple and “natural,” but not required. The freedom stems from the fact that we have only one condition coming from chiral scale invariance, but two LECs (C_v and C_s) with which we want to associate a^2 dependence. This lack of uniqueness is particularly apparent in the case $N_f = 2$, where the choice we made for Δ_{val} in Eq. (108) blows up. The problem is that, for $N_f = 2$, the change in the chiral logs

under change in chiral scale has no term that is proportional to m_x [see Eq. (107)]. In this case, one might wish to associate all the splittings with the sea quark. Of course, a redefinition like Eq. (111) or (112) is completely consistent for *any* choice of Δ_{sea} and Δ_{val} . Some redefinitions are “better” than others, however, if they result in a smaller C_a , and our choice does seem to work in the $N_f = 3$ case [9].

VIII. FINITE VOLUME CORRECTIONS

In computing the finite volume corrections, we assume that the lattice size in the time direction is larger than in the spatial directions, and indeed is large enough that it may be treated as infinite to a very good approximation. This is the case for the MILC lattices [10, 25]. We are thus interested in corrections due to finite *spatial* volume only. Furthermore, we assume that the heavy-light meson is at rest: $v^\mu = (1, 0, 0, 0)$. This is the case in all lattice simulations of the decay constant except those of “moving NRQCD” [39].

For integrals of the the form of Eqs. (93) and (94), which come from light-meson tadpole loops, the corrections are expressed in a convenient form in Ref. [12]. The results of these integrals appear in our final answers, Eqs. (95) through (106), in the terms multiplied by the “1” in the factor $1 + 3g_\pi^2$. For such terms, we merely need to make the replacements [12]

$$\ell(m^2) \rightarrow \ell_1(m^2) \equiv \ell(m^2) + m^2 \delta_1(mL) \quad [\text{finite volume, “1 terms”}] , \quad (114)$$

$$\tilde{\ell}(m^2) \rightarrow \tilde{\ell}_1(m^2) \equiv \tilde{\ell}(m^2) + \delta_3(mL) \quad [\text{finite volume, “1 terms”}] , \quad (115)$$

in Eqs. (91) and (92), where

$$\delta_1(mL) = \frac{4}{mL} \sum_{\vec{r} \neq 0} \frac{K_1(|\vec{r}|mL)}{|\vec{r}|} , \quad (116)$$

$$\delta_3(mL) = 2 \sum_{\vec{r} \neq 0} K_0(|\vec{r}|mL) , \quad (117)$$

with K_0 and K_1 the Bessel functions of imaginary argument.

The integrals that involve a heavy-light meson in the loop appear in our final answers in the terms multiplied by g_π^2 . For these integrals, Eqs. (89) and (90), the finite volume corrections can be determined by comparison with the corrections to Eqs. (93) and (94). To do this, we first perform the integrals over p^0 in both cases. This integration is not affected by finite volume corrections since the time extent of the lattices is taken to be infinite. We

obtain

$$\begin{aligned}
I_1(m^2) &= \int \frac{d^3p}{(2\pi)^3} \frac{1}{2\sqrt{\vec{p}^2 + m^2}} , \\
(g_{\mu\nu} - v_\mu v_\nu) \frac{\partial J^{\mu\nu}(m, \Delta)}{\partial \Delta} \Big|_{\Delta=0} &= - \int \frac{d^3p}{(2\pi)^3} \frac{\vec{p}^2}{2(\vec{p}^2 + m^2)^{3/2}} \\
&= - \left[I_1 + 2m^2 \frac{\partial I_1}{\partial m^2} \right] .
\end{aligned} \tag{118}$$

Since the finite volume correction to I_1 is given by Eq. (114), the correction to the heavy-light integral Eq. (89) is now determined *via* Eq. (118). Corrections to the integrals with double poles, Eqs. (94) and (90), follow by differentiation with respect to m^2 .

The prescription for finite volume corrections to the terms proportional to g_π^2 , is then

$$\ell(m^2) \rightarrow \ell_g(m^2) \equiv \ell(m^2) + \frac{m^2}{3} (\delta_1(mL) - 2\delta_3(mL)) \quad [\text{finite volume, “}g_\pi^2 \text{ terms”}] \tag{119}$$

$$\tilde{\ell}(m^2) \rightarrow \tilde{\ell}_g(m^2) \equiv \tilde{\ell}(m^2) + \delta_3(mL) - \frac{2}{3}\delta_5(mL) \quad [\text{finite volume, “}g_\pi^2 \text{ terms”}], \tag{120}$$

where

$$\delta_5(mL) \equiv -\frac{mL}{2} \delta'_3(mL) = mL \sum_{\vec{r} \neq 0} |\vec{r}| K_1(|\vec{r}|mL) , \tag{121}$$

with the prime denoting differentiation with respect to the argument. We have used $K'_0(z) = -K_1(z)$ and the relation $\delta_3(mL) = -\delta_1(mL) - (mL/2)\delta'_1(mL)$ [12].

We note that our procedure for finding finite volume corrections of heavy-light integrals works well only for the rather simple integrals needed here. It does not easily generalize to cases where the time extent is finite or where the variable Δ in Eq. (88) is nonzero. The latter case is particular important since it is needed for semileptonic form factors, as well as for $1/m_Q$ and $\mathcal{O}(m_q^2)$ corrections. The finite size effects for heavy-light χ Pt have been studied in much more generality in Ref. [40].

IX. REMARKS AND CONCLUSIONS

We have determined the heavy-light SXPT Lagrangian to NLO and used it to compute the heavy-light decay constant in partially quenched and full QCD. The heavy-light part of the chiral Lagrangian is identical at leading order to that in the continuum, aside from the extra taste degrees of freedom, which enter trivially. Thus taste-violating effects appear in the one-loop chiral logarithms only through the light-light meson sector, where the relevant

parameters have already been determined by simulations [10, 24, 25]. All the operators that appear at NLO in the heavy-light Lagrangian serve to produce only a term proportional to a^2 in the decay constants, and thus a single new low energy constant: c_a [Eqs. (95) through (106)] or C_a [Eq. (111) or (112)].

We emphasize that our calculations are based on the assumption that the staggered “fourth root trick” is valid. We further assume that this trick is correctly implemented on the chiral theory by insertions of factors of $1/4$ for each sea quark loop, where the loops are found by quark flow or replica analysis.

Our results most relevant at present are those for the $2+1$ case ($N_f = 3$ and $m_u = m_d$), since they apply to calculations using existing MILC configurations for the light dynamical quarks. These results have proved quite useful in reducing the errors from the chiral and continuum extrapolations in a lattice calculation of the leptonic decay constants of D and D_s mesons [9]. With the Lagrangian and the techniques presented here, one may also find expressions for the form factors for semileptonic heavy-to-light decays (such as $D \rightarrow K\ell\nu$ or $B \rightarrow \pi\ell\nu$) in SXPT. This has been done [18]; a detailed report of the calculation is in preparation [19]. The result has already been used in Ref. [8]. We also call attention to the work of Laiho [41], who has applied heavy-light SXPT to study heavy-to-heavy semileptonic decays, *e.g.*, $B \rightarrow D^*\ell\nu$.

A calculation of heavy-light B parameters in SXPT should also be straightforward. Although additional operators will contribute to analytic terms at NLO, the one-loop chiral logs themselves will not involve any new low energy constants beyond those already present in the continuum or in the one-loop SXPT for light-light mesons.

It is interesting to look at a simple example to see the effect of taste violations. In Fig. 6 we plot $f_{B_x}/f_{B_x}^{\text{LO}}$ for the partially quenched case with two degenerate sea quarks: Eq. (103) with $N_f = 2$. The solid line includes taste-breaking terms; while the dashed line shows the results with taste-breaking terms set to zero, *i.e.*, the continuum limit. We choose parameters from the coarse MILC data set ($a \approx 0.125\text{fm}$). The light meson masses and splittings, as well as the values of δ'_V and δ'_A , come from measurements of the light hadron spectrum and decay constants [10, 24, 25]. We have used the value $am_f = 0.010$ for the sea quark mass. We set $c_s = c_a = 0$ and choose c_v to give a slope at large mass similar to what is seen in simulations. The continuum result shows the characteristic divergence of partially quenched chiral perturbation theory as the valence mass is taken to zero for fixed sea mass.

The taste violations, however, completely wash out this behavior and give a result that looks rather linear. This is in fact an example of an infrared sensitive quantity for which the chiral and continuum limits in SXPT will not commute [42]. Fitting to and removing the effects of staggered taste violations is therefore crucial in controlling the systematic error associated with the extrapolation to the physical values of the light quark masses.

ACKNOWLEDGMENTS

We thank J. Bailey, B. Grinstein, A. Kronfeld, P. Mackenzie, S. Sharpe and our colleagues in the MILC collaboration for helpful discussions. This work was partially supported by the U.S. Department of Energy under grant number DE-FG02-91ER40628.

-
- [1] M. Wingate, Nucl. Phys. Proc. Suppl. **140**, 68 (2005) [arXiv:hep-lat/0410008].
 - [2] A. S. Kronfeld, Nucl. Phys. Proc. Suppl. **129**, 46 (2004) [arXiv:hep-lat/0310063].
 - [3] A. S. Kronfeld and S. M. Ryan, Phys. Lett. B **543**, 59 (2002) [arXiv:hep-ph/0206058]; N. Yamada, Nucl. Phys. **B** (Proc. Suppl.) **119** (2003), 93; N. Yamada *et al.*, Nucl. Phys. **B** (Proc. Suppl.) **106** (2002), 397 [arXiv:hep-lat/0110087].
 - [4] M. Wingate, *et al.*, Phys. Rev. D **67**, 054505 (2003) [arXiv:hep-lat/0211014].
 - [5] M. Wingate, C. Davies, A. Gray, E. Gulez, G. P. Lepage and J. Shigemitsu, Nucl. Phys. Proc. Suppl. **129**, 325 (2004) [arXiv:hep-lat/0309092].
 - [6] M. Wingate, C. T. H. Davies, A. Gray, G. P. Lepage and J. Shigemitsu, Phys. Rev. Lett. **92**, 162001 (2004) [arXiv:hep-ph/0311130].
 - [7] J. Shigemitsu, *et al.*, Nucl. Phys. Proc. Suppl. **129**, 331 (2004) [arXiv:hep-lat/0309039].
 - [8] C. Aubin *et al.* [Fermilab Lattice, MILC, and HPQCD Collaborations], Phys. Rev. Lett. **94**, 011601 (2005) [arXiv:hep-ph/0408306].
 - [9] C. Aubin *et al.* [Fermilab Lattice, MILC, and HPQCD Collaborations], Phys. Rev. Lett. **95**, 122002 (2005) [arXiv:hep-lat/0506030].
 - [10] C. Aubin *et al.* [MILC Collaboration], Phys. Rev. D **70**, 114501 (2004) [arXiv:hep-lat/0407028]; Phys. Rev. D **70**, 094505 (2004) [arXiv:hep-lat/0402030].
 - [11] W. Lee and S. Sharpe Phys. Rev. **D60**, 114503 (1999) [arXiv: hep-lat/9905023].
 - [12] C. Bernard, Phys. Rev. **D65**, 054031 (2001) [arXiv: hep-lat/0111051].

- [13] C. Aubin and C. Bernard, Phys. Rev. D **68**, 034014 (2003) [arXiv:hep-lat/0304014]; Phys. Rev. D **68**, 074011 (2003) [arXiv:hep-lat/0306026]; Nucl. Phys. **B** (Proc. Suppl.) **129-130C** (2004), 182 [arXiv:hep-lat/0308036].
- [14] E. Marinari, G. Parisi and C. Rebbi, Nucl. Phys. B **190**, 734 (1981).
- [15] S. Dürr and C. Hoelbling, Phys. Rev. **D69**, 034503 (2004) [arXiv:hep-lat/0311002]; E. Follana, A. Hart and C. T. H. Davies, Phys. Rev. Lett. **93**, 241601 (2004) [arXiv:hep-lat/0406010]; S. Dürr, C. Hoelbling and U. Wenger, Phys. Rev. **D70**, 094502 (2004) [arXiv:hep-lat/0406027]; S. Dürr and C. Hoelbling, Phys. Rev. **D71**, 054501 (2005) [arXiv:hep-lat/0411022]; F. Maresca and M. Peardon, arXiv:hep-lat/0411029; D. H. Adams, Phys. Rev. Lett. **92**, 162002 (2004) [arXiv:hep-lat/0312025] and arXiv:hep-lat/0411030; Y. Shamir, Phys. Rev. D **71**, 034509 (2005) [arXiv:hep-lat/0412014]; C. Bernard *et al.* [MILC Collaboration], Proceedings of Science, PoS(LAT2005)114, arXiv:hep-lat/0509176.
- [16] S. R. Sharpe, Phys. Rev. **D46**, 3146 (1992) [arXiv:hep-lat/9205020].
- [17] P. H. Damgaard and K. Splittorff, Phys. Rev. **D62**, 054509 (2000) [arXiv:hep-lat/0003017].
- [18] C. Aubin and C. Bernard, Nucl. Phys. **B** (Proc. Suppl.) **140** (2005), 491 [arXiv:hep-lat/0409027].
- [19] C. Aubin and C. Bernard, in preparation.
- [20] A. S. Kronfeld, Phys. Rev. D **62**, 014505 (2000) [arXiv:hep-lat/0002008].
- [21] C. Aubin, *et al.* [Fermilab Lattice, MILC and HPQCD Collaborations], *Heavy-to-light semileptonic decays of B and D mesons from three-flavor lattice QCD*, in preparation.
- [22] A. X. El-Khadra, A. S. Kronfeld and P. B. Mackenzie, Phys. Rev. D **55**, 3933 (1997) [arXiv:hep-lat/9604004].
- [23] G. P. Lepage and B. A. Thacker, Nucl. Phys. Proc. Suppl. **4**, 199 (1988); B. A. Thacker and G. P. Lepage, Phys. Rev. D **43**, 196 (1991).
- [24] C. Aubin *et al.* [MILC Collaboration], Nucl. Phys. **B** (Proc. Suppl.) **119** (2003), 233 [arXiv:hep-lat/0209066], Nucl. Phys. **B** (Proc. Suppl.) **129-130C** (2004), 227 [arXiv:hep-lat/0309088].
- [25] C. Bernard *et al.* [MILC Collaboration], Phys. Rev. **D64**, 054506, (2001) [arXiv:hep-lat/0104002]; Phys. Rev. **D62**, 034503, (2000) [arXiv:hep-lat/0002028]; Phys. Rev. **D61**, 111502(R), (2000) [arXiv:hep-lat/9912018].
- [26] M. Okamoto, private communication.

- [27] G. P. Lepage and P. B. Mackenzie, Phys. Rev. D **48**, 2250 (1993) [arXiv:hep-lat/9209022].
- [28] O. Bar, C. Bernard, G. Rupak and N. Shores, Phys. Rev. D **72**, 054502 (2005) [arXiv:hep-lat/0503009].
- [29] K. Orginos, D. Toussaint and R.L. Sugar, Phys. Rev. D **60** (1999) 054503 [arXiv:hep-lat/9903032].
- [30] S. R. Sharpe and R. S. Van de Water, Phys. Rev. D **71**, 114505 (2005) [arXiv:hep-lat/0409018].
- [31] G. Burdman and J. F. Donoghue, Phys. Lett. B **280**, 287 (1992); M. B. Wise, Phys. Rev. D **45**, 2188 (1992); T. M. Yan *et al.*, Phys. Rev. D **46**, 1148 (1992) [Erratum-ibid. D **55**, 5851 (1997)].
- [32] B. Grinstein, *et al.*, Nucl. Phys. B **380**, 369 (1992) [arXiv:hep-ph/9204207].
- [33] J. L. Goity, Phys. Rev. D **46**, 3929 (1992) [arXiv:hep-ph/9206230].
- [34] A. Manohar and M. Wise, *Heavy Quark Physics*, Cambridge University Press (2000) and references therein.
- [35] S. Sharpe and N. Shores, Phys. Rev. **D64**, 114510 (2001) [arXiv:hep-lat/0108003].
- [36] C. G. Boyd and B. Grinstein, Nucl. Phys. B **442**, 205 (1995) [arXiv:hep-ph/9402340].
- [37] I. W. Stewart, Nucl. Phys. B **529**, 62 (1998) [arXiv:hep-ph/9803227].
- [38] S. R. Sharpe and Y. Zhang, Phys. Rev. D **53**, 5125 (1996) [arXiv:hep-lat/9510037].
- [39] A. Dougall, K. M. Foley, C. T. H. Davies and G. P. Lepage, arXiv:hep-lat/0509108; K. M. Foley and G. P. Lepage, Nucl. Phys. Proc. Suppl. **119**, 635 (2003) [arXiv:hep-lat/0209135].
- [40] D. Arndt and C. J. D. Lin, Phys. Rev. D **70**, 014503 (2004) [arXiv:hep-lat/0403012].
- [41] J. Laiho, Proceedings of Science, PoS(LAT2005)221, arXiv:hep-lat/0510058.
- [42] C. Bernard, Phys. Rev. D **71**, 094020 (2005) [arXiv:hep-lat/0412030].




FIG. 1: Terms which contribute to the one-loop heavy-light meson decay constant, arising from wavefunction renormalization. The thick solid line is the heavy-light meson, and the dashed line is the light pseudoscalar (*i.e.*, “pion”). The solid square is the current insertion. The cross indicates a disconnected propagator, Eq. (75); a pion line without a cross is connected.




FIG. 2: Contributions to the one-loop heavy-light meson decay constant coming from corrections to the current insertion. Symbols have the same meaning as in Fig. 1.




FIG. 3: Terms which in principle arise at one loop in the decay constant but vanish since they are proportional to $v^\nu(g_{\mu\nu} - v_\mu v_\nu)$.




FIG. 4: Relevant vertices at the quark level. (a) is the g_π vertex. (b) is the second-order diagram coming from the current.




FIG. 5: Quark-flow diagrams that contribute to f_B , corresponding to Figs. 1 (left hand side) and 2 (right hand side). The dashed lines here are the light quarks while solid lines are the heavy quarks.




FIG. 6: The ratio of the NLO decay constant for two degenerate sea quarks, given by Eq. (103) with $N_f = 2$. The solid line is with the taste violations included, while the dashed line is the continuum limit.